arXiv:math/0703568v2 [math.RT] 16 Jun 2007

THE PRODUCT IN THE HOCHSCHILD COHOMOLOGY RING OF PREPROJECTIVE ALGEBRAS OF DYNKIN QUIVERS

CHING-HWA EU

Contents

1. Introduction 2
2. Preliminaries 3
2.1. Quivers and path algebras 3
2.2. The preprojective algebra 4
2.3. Graded spaces and Hilbert series 4
2.4. Root system parameters 4
3. Hochschild cohomology 5
4. Results about the product in the Hochschild cohomology ring of preprojective algebras for D- and E-quivers 6
5. Some basic facts about preprojective algebras 8
5.1. Labeling of quivers 8
5.2. Preprojective algebras by numbers 10
5.3. Frobenius algebras and Nakayama automorphism 10
5.4. The Schofield resolution 12
6. Basis and Hilbert series 13
6.1. $Q = D_{n+1}$ 13
6.2. $Q = E_6$ 13
6.3. $Q = E_7$ 14
6.4. $Q = E_8$ 17
7. $HH^0(A) = Z$ 20
7.1. $Q = D_{n+1}$ 20
7.2. $Q = E_6$ 21
7.3. $Q = E_7$ 22
7.4. $Q = E_8$ 23
8. $HH^1(A)$ 25
9. $HH^2(A)$ 26
9.1. $Q = D_{n+1}$, n even 26
9.2. $Q = E_6$ 27
10. $HH^3(A)$ 27
10.1. $Q = D_{n+1}$, n even 27
10.2. $Q = E_6$ 28
11. $HH^4(A)$ 28

11.1. $Q = D_{n+1}$, n odd 28
11.2. $Q = D_{n+1}$, n even 29
11.3. $Q = E_6$ 29
11.4. $Q = E_7$ 29
11.5. $Q = E_8$ 29
12. $HH^5(A)$ 29
12.1. $U^*[-2]$ 29
12.2. $Y^*[-h-2]$ 30
12.3. Result 34
13. $HH^6(A)$ 35
14. Products involving $HH^0(A) = Z$ 35
14.1. $HH^0(A) \times HH^0(A) \twoheadrightarrow HH^0(A)$ 35
14.2. $HH^0(A) \times HH^1(A) \twoheadrightarrow HH^1(A)$ 35
14.3. $HH^0(A) \times HH^i(A) \twoheadrightarrow HH^i(A)$, $i = 2$ or 3 36
14.4. $HH^0(A) \times HH^4(A) \twoheadrightarrow HH^4(A)$ 36
14.5. $HH^0(A) \times HH^5(A) \twoheadrightarrow HH^5(A)$ 36
15. Products involving $HH^1(A)$ 37
15.1. $HH^1(A) \times HH^1(A) \xrightarrow{0} HH^2(A)$ 37
15.2. $HH^1(A) \times HH^2(A) \twoheadrightarrow HH^3(A)$ 37
15.3. $HH^1(A) \times HH^3(A) \xrightarrow{0} HH^4(A)$ 40
15.4. $HH^1(A) \times HH^4(A) \to HH^5(A)$ 40
15.5. $HH^1(A) \times HH^5(A) \to HH^6(A)$ 41
16. Products involving $HH^2(A)$ 51
16.1. $HH^2(A) \times HH^3(A) \to HH^5(A)$ 51
16.2. $HH^2(A) \times HH^2(A) \to HH^4(A)$ 52
16.3. $HH^2(A) \times HH^4(A) \xrightarrow{0} HH^6(A)$ 53
16.4. $HH^2(A) \times HH^5(A) \xrightarrow{0} HH^7(A)$ 53
17. Products involving $HH^3(A)$ 53
17.1. $HH^3(A) \times HH^3(A) \xrightarrow{0} HH^6(A)$ 53
17.2. $HH^3(A) \times HH^4(A) \xrightarrow{0} HH^7(A)$ 53
17.3. $HH^3(A) \times HH^5(A) \xrightarrow{0} HH^8(A)$ 53
18. Products involving $HH^4(A)$ 53
18.1. $HH^4(A) \times HH^4(A) \xrightarrow{0} HH^8(A)$ 53
18.2. $HH^4(A) \times HH^5(A) \xrightarrow{0} HH^9(A)$ 54
19. $HH^5(A) \times HH^5(A) \to HH^{10}(A)$ 54
References 54

1. Introduction

In this paper, we compute the product structure of the Hochschild cohomology of preprojective algebras of quivers of type D and E over a field of characteristic zero. This is a continuation of [EE2] where the cohomology spaces together with the grading induced by the natural grading (all arrows have degree 1) were computed.

Together with the results in [ES2] where it was done for type A (over a field of any characteristic), this yields a complete description of the product in the Hochschild cohomology ring of preprojective algebras of ADE quivers over a field of characteristic zero.

We note that this description is essentially uniform (i.e. does not refer to particular Dynkin-diagrams), while the proof uses case-by-case arguments.

For our computation, the same complex as in [ES2] is used, namely the one which we get by applying the Hom-functor to the Schofield resolution (which is periodic with period 6) of the algebra.

To compute the cup product, we use the same method as in [ES2]: via the isomorphism $HH^i(A) \equiv \underline{Hom}(\Omega^i A, A)$ (where for an A-bimodule M we write ΩM for the kernel of its projective cover) we identify elements in $HH^i(A)$ with equivalence classes of maps $\Omega^i(A) \to A$. For $[f] \in HH^i(A)$ and $[g] \in HH^j(A)$, the product is $[f][g] := [f \circ \Omega^i g]$ in $HH^{i+j}(A)$. We compute all products $HH^i(A) \times HH^j(A) \to HH^{i+j}(A)$ for $0 \leq i \leq j \leq 5$. The remaining ones follow from the perodicity of the Schofield resolution and the graded commutativity of the multiplication. Some computations are similar to those in [ES2] for type A.

In the first part of the paper, we introduce a basis for each cohomology space explicitly (for each quiver). Then in the second part we compute the product in these bases. We use the results about the grading of the cohomology spaces from [EE2] to find the bases and the products.

Note that for connected non-Dynkin quivers, the Hochschild cohomology and its product structure were already calculated in [CBEG] where the situation is much easier because the homological dimension of the preprojective algebra is 2.

Acknowledgements. C. Eu wants to thank his advisor P. Etingof and T. Schedler for useful discussions. This work is partially supported by the NSF grant DMS-0504847.

2. Preliminaries

2.1. **Quivers and path algebras.** Let Q be a quiver of ADE type with vertex set I and $|I| = r$. We write $a \in Q$ to say that a is an arrow in Q.

We define Q^* to be the quiver obtained from Q by reversing all of its arrows. We call $\bar{Q} = Q \cup Q^*$ the *double* of Q.

Let C be the adjacency matrix corresponding to the quiver $\bar{Q}$.

The concatenation of arrows generate the *nontrivial paths* inside the quiver $\bar{Q}$. We define e_i, $i \in I$ to be the *trivial path* which starts and ends at i. The *path algebra* $P_{\bar{Q}} = \mathbb{C}\bar{Q}$ of $\bar{Q}$ over $\mathbb{C}$ is the $\mathbb{C}$-algebra with basis the paths in $\bar{Q}$ and the product xy of two paths x and y to be their concatenation if they are compatible and 0 if not. We define the *Lie bracket* $[x,y]=xy-yx$.

Let $R=\oplus_{i\in I}\mathbb{C}e_i$. Then R is a commutative semisimple algebra, and P_Q is naturally an R-bimodule.

2.2. **The preprojective algebra.** Given a quiver Q, we define the *preprojective algebra* Π_Q to be the quotient of the path algebra $P_{\bar{Q}}$ by the relation $\sum\limits_{a\in Q}[a,a^*]=0$.

Given a path x, we write x^* for the path obtained from x by reversing all arrows.

From now on, we write $A=\Pi_Q$.

2.3. **Graded spaces and Hilbert series.** Let $M=\oplus_{d\geq 0}M(d)$ be a $\mathbb{Z}_+$-graded vector space, with finite dimensional homogeneous subspaces. We denote by $M[n]$ the same space with grading shifted by n. The graded dual space M^* is defined by the formula $M^*(n)=M(-n)^*$.

Definition 2.3.1. (The Hilbert series of vector spaces)
We define the Hilbert series $h_M(t)$ *to be the series*

$$h_M(t)=\sum_{d=0}^{\infty}\dim M(d)t^d.$$

Definition 2.3.2. (The Hilbert series of bimodules)
Let $M=\oplus_{d\geq 0}M(d)$ be a $\mathbb{Z}_+$-graded bimodule over the ring R, so we can write $M=\oplus M_{i,j}$. We define the Hilbert series $H_M(t)$ *to be a matrix valued series with the entries*

$$H_M(t)_{i,j}=\sum_{d=0}^{\infty}\dim\ M(d)_{i,j}t^d.$$

2.4. **Root system parameters.** Let w_0 be the longest element of the Weyl group W of Q. Then we define ν to be the involution of I, such that $w_0(\alpha_i)=-\alpha_{\nu(i)}$ (where α_i is the simple root corresponding to $i\in I$). It turns out that $\eta(e_i)=e_{\nu(i)}$ ([S]; see [ES2]).

Let m_i, $i=1,...,r$, be the exponents of the root system attached to Q, enumerated in the increasing order. Let $h=m_r+1$ be the Coxeter number of Q.

Let P be the permutation matrix corresponding to the involution ν. Let $r_+=\dim\ker(P-1)$ and $r_-=\dim\ker(P+1)$. Thus, r_- is half the number of vertices which are not fixed by ν, and $r_+=r-r_-$.

A is finite dimensional, and the following Hilbert series is known from [MOV, Theorem 2.3.]:

$$H_A(t)=(1+Pt^h)(1-Ct+t^2)^{-1}. \tag{2.4.1}$$

We see that the top degree of A is $h-2$, and for the top degree A^{top} part we get the following decomposition in 1-dimensional submodules:

$$A^{top} = A(h-2) = \bigoplus_{i\in I} e_i A(h-2) e_{\nu(i)} \tag{2.4.2}$$

3. HOCHSCHILD COHOMOLOGY

The Hochschild cohomology spaces of A were computed in [EE2]. We recall the results:

Definition 3.0.3. *We define the spaces*

$$\begin{aligned} U &= \oplus_{d<h-2} HH^0(A)(d)[2],\\ L &= HH^0(A)(h-2),\\ K &= HH^2(A)[2],\\ Y &= HH^6(A)(-h-2). \end{aligned}$$

Theorem 3.0.4. (1) *U has the following Hilbert series:*

$$h_U(t) = \sum_{\substack{i=1\\ m_i<\frac{h}{2}}}^{r} t^{2m_i}. \tag{3.0.5}$$

(2) *We have natural isomorphisms*

$$K \equiv \ker(P+1),$$
$$L \equiv \ker(P-1),$$

and

$$\dim Y = r_+ - r_- - \#\{i : m_i = \frac{h}{2}\}.$$

Theorem 3.0.6. *For the Hochschild cohomology spaces, we have the following natural isomorphisms:*

$$\begin{aligned} &HH^0(A) = U[-2] \oplus L[h-2],\\ &HH^1(A) = U[-2],\\ &HH^2(A) = K[-2],\\ &HH^3(A) = K^*[-2],\\ &HH^4(A) = U^*[-2],\\ &HH^5(A) = U^*[-2] \oplus Y^*[-h-2],\\ &HH^6(A) = U[-2h-2] \oplus Y[-h-2], \end{aligned}$$

and $HH^{6n+i}(A) = HH^i(A)[-2nh]\,\forall i \geq 1.$

Corollary 3.0.7. *The center* $Z = HH^0(A)$ *of* A *has Hilbert series*

$$h_Z(t) = \sum_{\substack{i=1 \\ m_i < \frac{h}{2}}}^{r} t^{2m_i-2} + r_+ t^{h-2}.$$

4. RESULTS ABOUT THE PRODUCT IN THE HOCHSCHILD COHOMOLOGY RING OF PREPROJECTIVE ALGEBRAS FOR D- AND E-QUIVERS

Let $(U[-2])_+$ be the positive degree part of $U[-2]$ (which lies in non-negative degrees).

We have a decomposition $HH^0(A) = \mathbb{C} \oplus (U[-2])_+ \oplus L[-h-2]$ where we have the natural identification $(U[-2])(0) = \mathbb{C}$.

We give a brief description of the product structure in $HH^*(A)$ which will be computed in this paper. Since the product $HH^i(A) \times HH^j(A) \to HH^{i+j}(A)$ is graded-commutative, we can assume $i \leq j$ here.

Let $z_0 = 1 \in \mathbb{C} \subset U[-2] \subset HH^0(A)$ (in lowest degree 0),
θ_0 the corresponding element in $HH^1(A)$ (in lowest degree 0),
ψ_0 the dual element of z_0 in $U^*[-2] \subset HH^5(A)$ (in highest degree -4), i.e. $\psi_0(z_0) = 1$,
ζ_0 the corresponding element in $U^*[-2] \subset HH^4(A)$ (in highest degree -4), that is the dual element of θ_0, $\zeta_0(\theta_0) = 1$,
$\varphi_0 : HH^0(A) \to HH^6(A)$ the natural quotient map (which induces the natural isomorphism $U[-2] \to U[-2h-2]$) and
ϕ the quotient map $L \to Y$ induced by φ_0.

Theorem 4.0.8. *(The product structure in* $HH^*(A)$ *for quivers of type D and E)*

1. *The multiplication by* $\varphi_0(z_0)$ *induces the natural isomorphisms* $\varphi_i : HH^i(A) \to HH^{i+6}(A)$ $\forall i \geq 1$ *and the natural quotient map* φ_0. *Therefore, it is enough to compute products* $HH^i(A) \times HH^j(A) \to HH^{i+j}(A)$ *with* $0 \leq i \leq j \leq 5$.
2. *The* $HH^0(A)$*-action on* $HH^i(A)$*:*
   - (a) *(*$(U[-2])_+$*-action)*
     The action of $(U[-2])_+$ *on* $U[-2] \subset HH^1(A)$ *corresponds to the multiplication*

$$\begin{aligned} (U[-2])_+ \times U[-2] &\to U[-2], \\ (u, v) &\mapsto u \cdot v \end{aligned}$$

in $HH^0(A)$*, projected on* $U[-2] \subset HH^0(A)$.
$(U[-2])_+$ *acts on* $U^*[-2] = HH^4(A)$ *and* $U^*[-2] \subset HH^5(A)$ *the following way:*

$$\begin{aligned} (U[-2])_+ \times U^*[-2] &\to U^*[-2], \\ (u, f) &\mapsto u \circ f, \end{aligned}$$

where $(u \circ f)(v) = f(uv)$.
$(U[-2])_+$ *acts by zero on* $L[h-2] \subset HH^0(A)$, $HH^2(A)$, $HH^3(A)$ *and* $Y^*[-h-2] \subset HH^5(A)$.

(b) *(*$L[h-2]$*-action)*
$L[h-2]$ *acts by zero on* $HH^i(A)$, $1 \leq i \leq 4$, *and on* $U^*[-2] \subset HH^5(A)$.
The $L[h-2]$*-action on* $HH^5(A)$ *restricts to*

$$\begin{array}{rcl} L[h-2] \times Y^*[-h-2] & \to & U^*[-2], \\ (a, y) & \mapsto & y(\phi(a))\psi_0. \end{array}$$

(3) *(Zero products)*
All products $HH^i(A) \times HH^j(A) \to HH^{i+j}$, $1 \leq i \leq j \leq 5$, *where* $i + j \geq 6$ *or* i, j *are both odd are zero except the pairings*

$$HH^1(A) \times HH^5(A) \to HH^6(A)$$

and

$$HH^5(A) \times HH^5(A) \to HH^{10}(A).$$

(4) *(*$HH^1(A)$*-products)*

(a) *The multiplication*

$$HH^1(A) \times HH^4(A) = U[-2] \times U^*[-2] \to HH^5(A)$$

is the same one as the restriction of

$$HH^0(A) \times HH^5(A) \to HH^5(A)$$

on $U[-2] \times U^*[-2]$.

(b) *The multiplication of the subspace* $U[-2]_+ \subset HH^1(A)$ *with* $HH^i(A)$ *where* $i = 2, 5$ *is zero.*

(c) *The multiplication by* θ_0 *induces a symmetric isomorphism*

$$\alpha : HH^2(A) = K[-2] \to K^*[-2] = HH^3(A).$$

On $HH^5(A)$, *it induces a skew-symmetric isomorphism*

$$\beta : Y^*[-h-2] \to Y[-h-2] \subset HH^6(A),$$

and acts by zero on $U^*[-2] \subset HH^5(A)$. α *and* β *will be given by explicit matrices* M_α *amd* M_β *later.*

(5) *(*$HH^2(A)$*-products)*

$$\begin{array}{rcl} HH^2(A) \times HH^2(A) & \to & HH^4(A), \\ (a, b) & \mapsto & \langle -, - \rangle \zeta_0 \end{array}$$

is given by $\langle -, - \rangle = \alpha$ *where* α *is regarded as a symmetric bilinear form.*

$HH^2(A) \times HH^3(A) \to HH^5(A)$ is the multiplication

$$\begin{array}{rcl} K[-2] \times K^*[-2] & \to & HH^5(A), \\ (a, y) & \mapsto & y(a)\psi_0. \end{array}$$

(6) *($HH^5(A) \times HH^5(A) \to HH^{10}(A)$)*
The restriction of this product to

$$\begin{array}{rcl} Y^*[-h-2] \times Y^*[-h-2] & \to & HH^{10}(A), \\ (a, b) & \mapsto & \Omega(-,-)\varphi_4(\zeta_0) \end{array}$$

is given by $\Omega(-,-) = -\beta$ where β is regarded as a skew-symmetric bilinear form.

The multiplication of the subspace $U^*[-2] \subset HH^5(A)$ with $HH^5(A)$ is zero.

5. Some basic facts about preprojective algebras

5.1. **Labeling of quivers.** From now on, we use the following labellings for the different types of quivers:



Figure 1. D_{n+1}-quiver

5.1.1. $Q = D_{n+1}$. A is the path algebra modulo the relations

$$\begin{array}{rcl} a^*_1 a_1 & = & 0, \\ a^*_{i+1} a_{i+1} & = & a_i a^*_i, \quad 1 \leq n-3 \\ a^*_{n-1} a_{n-1} + a^*_n a_n & = & a_{n-2} a^*_{n-2}. \end{array}$$

5.1.2. $Q = E_6$. A is the path algebra modulo the relations

$$\begin{array}{rcl} a_1 a^*_1 = a_4 a^*_4 = a_5 a^*_5 & = & 0, \\ a^*_1 a_1 & = & a_2 a^*_2, \\ a^*_4 a_4 & = & a_3 a^*_3, \\ a^*_2 a_2 + a^*_3 a_3 + a_5 a^*_5 & = & 0. \end{array}$$



FIGURE 2. E_6-quiver



FIGURE 3. E_7-quiver

5.1.3. $Q = E_7$. A is the path algebra modulo the relations

$$\begin{aligned} a_1a_1^* = a_5a_5^* = a_6a_6^* &= 0, \\ a_1^*a_1 &= a_2a_2^*, \\ a_2^*a_2 &= a_3a_3^*, \\ a_5^*a_5 &= a_4a_4^*, \\ a_3^*a_3 + a_4^*a_4 + a_6a_6^* &= 0. \end{aligned}$$



FIGURE 4. E_8-quiver

A is the path algebra modulo the relations

$$\begin{aligned} a_0a_0^* = a_5a_5^* = a_6a_6^* &= 0, \\ a_0^*a_0 &= a_1a_1^*, \\ a_1^*a_1 &= a_2a_2^*, \\ a_2^*a_2 &= a_3a_3^*, \\ a_5^*a_5 &= a_4a_4^*, \\ a_3^*a_3 + a_4^*a_4 + a_6a_6^* &= 0. \end{aligned}$$

5.2. **Preprojective algebras by numbers.** We summarize useful numbers associated to preprojective algebras, by quiver:

Q		exponents m_i	h	$\deg A^{top}$	degrees $HH^0(A)$
D_{n+1}	n odd	$1,3,\dots,2n-1,n$	$2n$	$2n-2$	$0,4,\dots,2n-6,2n-2$
	n even				$0,4,\dots 2n-4,2n-2$
E_6		$1,4,5,7,8,11$	12	10	$0,6,8,10$
E_7		$1,5,7,9,11,13,17$	18	16	$0,8,12,16$
E_8		$1,7,11,13,17,19,23,29$	30	28	$0,12,20,24,28$

We see that for quivers of type D and E, the degrees of the space U (which are $2m_i$, $m_i < \frac{h}{2}$) are even and range from 0 to $h-2$.

We get the following degree ranges for the Hochschild cohomology:

$$\begin{array}{lll} HH^0(A) & = U[-2] \oplus L[h-2] & 0 \leq \deg HH^0(A) \leq h-2 \\ HH^1(A) & = U[-2] & 0 \leq \deg HH^1(A) \leq h-4 \\ HH^2(A) & = K[-2] & \deg HH^2(A) = -2 \\ HH^3(A) & = K[-2] & \deg HH^3(A) = -2 \\ HH^4(A) & = U^*[-2] & -h \leq \deg HH^4(A) \leq -4 \\ HH^5(A) & = U^*[-2] \oplus Y^*[-h-2] & -h-2 \leq \deg HH^5(A) \leq -4 \\ HH^6(A) & = U[-2h-2] \oplus Y[-h-2] & -2h \leq \deg HH^6(A) \leq -h-2 \end{array}$$

5.3. Frobenius algebras and Nakayama automorphism.

Definition 5.3.1. *Let $\mathcal{A}$ be a finite dimensional unital $\mathbb{C}-$algebra. We call it Frobenius if there is a linear function $f : \mathcal{A} \to \mathbb{C}$, such that the form $(x, y) := f(xy)$ is nondegenerate, or, equivalently, if there exists an isomorphism $\phi : \mathcal{A} \overset{\simeq}{\to} \mathcal{A}^*$ of left $\mathcal{A}-$modules: given f, we can define $\phi(a)(b) = f(ba)$, and given ϕ, we define $f = \phi(1)$.*

Remark 5.3.2. If $\tilde{f}$ is another linear function satisfying the same properties as f from above, then $\tilde{f}(x) = f(xa)$ for some invertible $a \in \mathcal{A}$. Indeed, we define the form $\{a, b\} = \tilde{f}(ab)$. Then $\{-, 1\} \in \mathcal{A}^*$, so there is an $a \in \mathcal{A}$, such that $\phi(a) = \{-, 1\}$. Then $\tilde{f}(x) = \{x, 1\} = \phi(a)(x) = f(xa)$.

Definition 5.3.3. *Given a Frobenius algebra $\mathcal{A}$ (with a function f inducing a bilinear form $(-, -)$ from above), the automorphism $\eta : \mathcal{A} \to \mathcal{A}$ defined by the equation $(x, y) = (y, \eta(x))$ is called the* Nakayama automorphism *(corresponding to f).*

Remark 5.3.4. We note that the freedom in choosing f implies that η is uniquely determined up to an inner automorphism. Indeed, let $\tilde{f}(x) = f(xa)$ and define the bilinear form $\{a, b\} = \tilde{f}(ab)$. Then

$$\begin{aligned}\{x, y\} = \tilde{f}(xy) = f(xya) = (x, ya) = (ya, \eta(x)) = f(ya\eta(x)a^{-1}a)\\ = (y, a\eta(x)a^{-1}).\end{aligned}$$

It is known that A is a Frobenius algebra (see e.g. [ES2],[MOV]).

The linear function $f : A \to \mathbb{C}$ is zero in the non-top degree part of A. It maps a top degree element $\omega_i \in e_i A^{top} e_{\nu(i)}$ to 1. It is uniquely determined by the choice of one of these ω_i and a Nakayama automorphism.

For each quiver, we define a Nakayama automorphism and make a choice of one $\omega_i \in e_i A^{top} e_{\nu(i)}$:

5.3.1. *$Q = D_{n+1}$, n odd.* We define η by

$$\eta(a_i) = -a_i, \tag{5.3.5}$$

$$\eta(a_i^*) = a_i^*, \tag{5.3.6}$$

(5.3.7)

and

$$\omega_1 = a_1^* \ldots a_{n-2}^* a_{n-1}^* a_{n-1} a_{n-2} \ldots a_1. \tag{5.3.8}$$

5.3.2. *$Q = D_{n+1}$, n even.* We define η by

$$\forall i \leq n-2: \quad \eta(a_i) = -a_i, \tag{5.3.9}$$

$$\forall i \leq n-2: \quad \eta(a_i^*) = a_i^*, \tag{5.3.10}$$

$$\eta(a_{n-1}) = -a_n, \tag{5.3.11}$$

$$\eta(a_{n-1}^*) = a_n^*, \tag{5.3.12}$$

$$\eta(a_n) = -a_{n-1}, \tag{5.3.13}$$

$$\eta(a_n^*) = a_{n-1}^*, \tag{5.3.14}$$

$$\omega_1 = a_1^* \ldots a_{n-2}^* a_{n-1}^* a_{n-1} a_{n-2} \ldots a_1. \tag{5.3.15}$$

5.3.3. *$Q = E_6$.* We define η by

$$\eta(a_1) = -a_4, \tag{5.3.16}$$

$$\eta(a_1^*) = a_4^*, \tag{5.3.17}$$

$$\eta(a_2) = -a_3, \tag{5.3.18}$$

$$\eta(a_2^*) = a_3^*, \tag{5.3.19}$$

$$\eta(a_5) = -a_5, \tag{5.3.20}$$

$$\eta(a_5^*) = a_5^*, \tag{5.3.21}$$

and

$$\omega_3 = a_3^* a_3 (a_2^* a_2 a_3^* a_3)^2. \tag{5.3.22}$$

5.3.4. $Q = E_7$. We define η by

$$\eta(a_i) = -a_i, \tag{5.3.23}$$

$$\eta(a_i^*) = a_i^*, \tag{5.3.24}$$

and

$$\omega_4 = (a_4^* a_4 a_3^* a_3)^4. \tag{5.3.25}$$

5.3.5. $Q = E_8$. We define η by

$$\eta(a_i) = -a_i, \tag{5.3.26}$$

$$\eta(a_i^*) = a_i^*, \tag{5.3.27}$$

and

$$\omega_4 = (a_4^* a_4 a_3^* a_3)^7 \tag{5.3.28}$$

5.4. **The Schofield resolution.** We recall the Schofield resolution of A from [S].

Define the $A-$bimodule $\mathcal{N}$ obtained from A by twisting the right action by η, i.e., $\mathcal{N} = A$ as a vector space, and $\forall a, b \in A, x \in \mathcal{N} : a \cdot x \cdot b = ax\eta(b)$. Introduce the notation $\epsilon_a = 1$ if $a \in Q$, $\epsilon_a = -1$ if $a \in Q^*$. Let x_i be a homogeneous basis of A and x_i^* the dual basis under the form attached to the Frobenius algebra A. Let V be the bimodule spanned by the edges of $\bar{Q}$.

We start with the following exact sequence:

$$0 \to \mathcal{N}[h] \overset{i}{\to} P_2 \overset{d_2}{\to} P_1 \overset{d_1}{\to} P_0 \overset{d_0}{\to} A \to 0,$$

where $P_2 = A \otimes_R A[2]$, $P_1 = A \otimes_R V \otimes_R A$, $P_0 = A \otimes_R A$,

$$\begin{aligned} d_0(x \otimes y) &= xy, \\ d_1(x \otimes v \otimes y) &= xv \otimes y - x \otimes vy, \\ d_2(z \otimes t) &= \sum_{a \in \bar{Q}} \epsilon_a za \otimes a^* \otimes t + \sum_{a \in \bar{Q}} \epsilon_a z \otimes a \otimes a^* t, \\ i(a) &= a \sum x_i \otimes x_i^*. \end{aligned}$$

Since $\eta^2 = 1$, we can make a canonical identification $A = \mathcal{N} \otimes_A \mathcal{N}$ (via $x \mapsto x \otimes 1$), so by tensoring the above exact sequence with $\mathcal{N}$, connecting with the original exact sequence and repeating this process, we get the Schofield resolution

$$\ldots \to P_6 \overset{d_6}{\to} P_5 \overset{d_5}{\to} P_4 \overset{d_4}{\to} P_3 \overset{d_3}{\to} P_2 \overset{d_2}{\to} P_1 \overset{d_1}{\to} P_0 \overset{d_0}{\to} A \to 0,$$

with

$$P_{i+3} = (P_i \otimes_R \mathcal{N})[h].$$

We will work with the Hochschild cohomology complex obtained from this resolution, which is given explicitly in [EE2, Subsection 4.5.].

6. BASIS AND HILBERT SERIES

We need to work with the Hilbert series and with an explicit basis of A. We do this for each type of quiver separately.

We write B for a set of all basis elements of A, $B_{i,-}$ for a basis of e_iA, $B_{-,j}$ for a basis of Ae_j, $B_{i,j}$ for a basis of e_iAe_j and $B_{i,j}(d)$ for a basis of $e_iAe_j(d)$.

6.1. $Q = D_{n+1}$**.** A basis of A is given by the following elements:

For $\underline{k, j \leq n-1}$:

$$\begin{array}{rcl}
B_{k,n} & = & \{(a_{k-1}a_{k-1}^*)^l a_k^* \cdots a_{n-2}^* a_{n-1}^* | 0 \leq l \leq k-1\}, \\
B_{k,n+1} & = & \{(a_{k-1}a_{k-1}^*)^l a_k^* \cdots a_{n-2}^* a_n^* | 0 \leq l \leq k-1\}, \\
B_{n,n} & = & \{(a_{n-1}a_n^* a_n a_{n-1}^*)^l | 0 \leq l \leq \left\{ \begin{array}{ll} \frac{n-1}{2} & n\, odd, \\ \frac{n-2}{2} & n\, even \end{array} \right. \}, \\
B_{n+1,n+1} & = & \{(a_n a_{n-1}^* a_{n-1} a_n^*)^l | 0 \leq l \leq \left\{ \begin{array}{ll} \frac{n-1}{2} & n\, odd, \\ \frac{n-2}{2} & n\, even \end{array} \right. \}, \\
B_{n+1,n} & = & \{a_n a_{n-1}^* (a_{n-1} a_n^* a_n a_{n-1}^*)^l | 0 \leq l \leq \left\{ \begin{array}{ll} \frac{n-3}{2} & n\, odd, \\ \frac{n-2}{2} & n\, even \end{array} \right. \}, \\
B_{n,n+1} & = & \{a_{n-1} a_n^* (a_n a_{n-1}^* a_{n-1} a_n^*)^l | 0 \leq l \leq \left\{ \begin{array}{ll} \frac{n-3}{2} & n\, odd, \\ \frac{n-2}{2} & n\, even \end{array} \right. \}, \\
B_{n,j} & = & \{a_{n-1} a_{n-2} \cdots a_j (a_{j-1} a_{j-1}^*)^l | 0 \leq l \leq j-1\}, \\
B_{n+1,j} & = & \{a_n a_{n-2} \cdots a_j (a_{j-1} a_{j-1}^*)^l | 0 \leq l \leq j-1\}.
\end{array}$$

For $\underline{k \leq j \leq n-1}$,

$$\begin{array}{rcl}
B_{k,j} & = & \{(a_{k-1}a_{k-1}^*)^l a_k^* \cdots a_{j-1}^* | 0 \leq l \leq \min\{k-1, n-j-1\}\} \cup \\
& & \{(a_{k-1}a_{k-1}^*)^l a_k^* \cdots a_{n-1}^* a_{n-1} a_{n-2} a_j | 0 \leq l \leq k-1\} \cup \\
& & \{(a_{k-1}a_{k-1}^*)^l a_k^* \cdots a_n^* a_n a_{n-2} a_j | 0 \leq l \leq k-1+j-n\}.
\end{array}$$

For $\underline{j < k \leq n-1}$,

$$\begin{array}{rcl}
B_{k,j} & = & \{a_{k-1} \cdots a_j (a_j^* a_j)^l | 0 \leq l \leq \min\{n-k-1, j-1\}\} \cup \\
& & \{a_k^* \cdots a_{n-2}^* a_{n-1}^* a_{n-1} a_{n-2} \cdots a_j (a_j^* a_j)^l | 0 \leq l \leq j-1\} \cup \\
& & \{a_k^* \cdots a_{n-2}^* a_n^* a_n a_{n-2} \cdots a_j (a_j^* a_j)^l | 0 \leq l \leq j-1+k-n\}.
\end{array}$$

6.2. $Q = E_6$**.** We give the columns of the Hilbert series $H_A(t)$ which can be calculated from 2.4.1:

$$(H_A(t)_{i,1})_{1 \leq i \leq 6} = \left(\begin{array}{l} 1+t^6 \\ t+t^5+t^7 \\ t^2+t^4+t^6+t^8 \\ t^3+t^5+t^9 \\ t^4+t^{10} \\ t^3+t^7 \end{array} \right),$$

$$(H_A(t)_{i,2})_{1\le i\le 6} = \begin{pmatrix} t+t^5+t^7 \\ 1+t^2+t^4+2t^6+t^8 \\ t+2t^3+2t^5+2t^7+t^9 \\ t^2+2t^4+t^6+t^8+t^{10} \\ t^3+t^5+t^9 \\ t^2+t^4+t^6+t^8 \end{pmatrix},$$

$$(H_A(t)_{i,3})_{1\le i\le 6} = \begin{pmatrix} t^2+t^4+t^6+t^8 \\ t+2t^3+2t^5+2t^7+t^9 \\ 1+2t^2+3t^4+3t^6+2t^8+t^{10} \\ t+2t^3+2t^5+2t^7+t^9 \\ t^2+t^4+t^6+t^8 \\ t+t^3+2t^5+t^7+t^9 \end{pmatrix},$$

$$(H_A(t)_{i,4})_{1\le i\le 6} = \begin{pmatrix} t^3+t^5+t^9 \\ t^2+2t^4+t^6+t^8+t^{10} \\ t+2t^3+2t^5+2t^7+t^9 \\ 1+t^2+t^4+2t^6+t^8 \\ t+t^5+t^7 \\ t^2+t^4+t^6+t^8 \end{pmatrix},$$

$$(H_A(t)_{i,5})_{1\le i\le 6} = \begin{pmatrix} t^4+t^{10} \\ t^3+t^5+t^9 \\ t^2+t^4+t^6+t^8 \\ t+t^5+t^7 \\ 1+t^6 \\ t^3+t^7 \end{pmatrix},$$

$$(H_A(t)_{i,6})_{1\le i\le 6} = \begin{pmatrix} t^3+t^7 \\ t^2+t^4+t^6+t^8 \\ t+t^3+2t^5+t^7+t^9 \\ t^2+t^4+t^6+t^8 \\ t^3+t^7 \\ 1+t^4+t^6+t^{10} \end{pmatrix}.$$

6.3. $Q = E_7$**.** We give the columns of the Hilbert series matrix $H_A(t)$ of A which can be calculated from 2.4.1:

$$(H_A(t)_{i,1})_{1\le i\le 7} = \begin{pmatrix} 1+t^8+t^{16} \\ t+t^7+t^9+t^{15} \\ t^2+t^6+t^8+t^{10}+t^{14} \\ t^3+t^5+t^7+t^9+t^{11}+t^{13} \\ t^4+t^6+t^{10}+t^{12} \\ t^5+t^{11} \\ t^4+t^8+t^{12} \end{pmatrix},$$

$$(H_A(t)_{i,2})_{1\leq i\leq 7} = \begin{pmatrix} t+t^7+t^9+t^{15} \\ 1+t^2+t^6+2t^8+t^{10}+t^{14}+t^{16} \\ t+t^3+t^5+2t^7+2t^9+t^{11}+t^{13}+t^{15} \\ t^2+2t^4+2t^6+2t^8+2t^{10}+2t^{12}+t^{14} \\ t^3+2t^5+t^7+t^9+2t^{11}+t^{13} \\ t^4+t^6+t^{10}+t^{12} \\ t^3+t^5+t^7+t^9+t^{11}+t^{13} \end{pmatrix},$$

$$(H_A(t)_{i,3})_{1\leq i\leq 7} = \begin{pmatrix} t^2+t^6+t^8+t^{10}+t^{14} \\ t+t^3+t^5+2t^7+2t^9+t^{11}+t^{13}+t^{15} \\ 1+t^2+2t^4+3t^6+3t^8+2t^{10}+2t^{12}+t^{14}+t^{16} \\ 1+2t^2+3t^4+4t^6+4t^8+4t^{10}+3t^{12}+2t^{14}+t^{16} \\ t+2t^3+2t^5+3t^7+3t^9+2t^{11}+2t^{13}+t^{15} \\ t^2+t^4+t^6+2t^8+t^{10}+t^{12}+t^{14} \\ t+t^3+2t^5+2t^7+2t^9+2t^{11}+t^{13}+t^{15} \end{pmatrix}$$

$$(H_A(t)_{i,4})_{1\leq i\leq 7} = \begin{pmatrix} t^3+t^5+t^7+t^9+t^{11}+t^{13} \\ t^2+2t^4+2t^6+2t^8+2t^{10}+2t^{12}+t^{14} \\ t+2t^3+3t^5+3t^7+3t^9+3t^{11}+2t^{13}+t^{15} \\ 1+2t^2+3t^4+4t^6+4t^8+4t^{10}+3t^{12}+2t^{14}+t^{16} \\ t+2t^3+2t^5+3t^7+3t^9+2t^{11}+2t^{13}+t^{15} \\ t^2+t^4+t^6+2t^8+t^{10}+t^{12}+t^{14} \\ t+t^3+2t^5+2t^7+2t^9+2t^{11}+t^{13}+t^{15} \end{pmatrix},$$

$$(H_A(t)_{i,5})_{1\leq i\leq 7} = \begin{pmatrix} t^4+t^6+t^{10}+t^{12} \\ t^3+2t^5+t^7+t^9+2t^{11}+t^{13} \\ t^2+2t^4+2t^6+2t^8+2t^{10}+2t^{12}+t^{14} \\ t+2t^3+2t^5+3t^7+3t^9+2t^{11}+2t^{13}+t^{15} \\ 1+t^2+t^4+2t^6+2t^8+2t^{10}+t^{12}+t^{14}+t^{16} \\ t+t^5+t^7+t^9+t^{11}+t^{15} \\ t^2+t^4+t^6+2t^8+t^{10}+t^{12}+t^{14} \end{pmatrix},$$

$$(H_A(t)_{i,6})_{1\leq i\leq 7} = \begin{pmatrix} t^5+t^{11} \\ t^4+t^6+t^{10}+t^{12} \\ t^3+t^5+t^7+t^9+t^{11}+t^{13} \\ t^2+t^4+t^6+2t^8+t^{10}+t^{12}+t^{14} \\ t+t^5+t^7+t^9+t^{11}+t^{15} \\ 1+t^6+t^{10}+t^{16} \\ t^3+t^7+t^9+t^{13} \end{pmatrix},$$

$$(H_A(t)_{i,7})_{1\leq i\leq 7} = \begin{pmatrix} t^4+t^8+t^{12} \\ t^3+t^5+t^7+t^9+t^{11}+t^{13} \\ t^2+t^4+2t^6+t^8+2t^{10}+t^{12}+t^{14} \\ t+t^3+2t^5+2t^7+2t^9+2t^{11}+t^{13}+t^{15} \\ t^2+t^4+t^6+2t^8+t^{10}+t^{12}+t^{14} \\ t^3+t^7+t^9+t^{13} \\ 1+t^4+t^6+t^8+t^{10}+t^{12}+t^{16} \end{pmatrix}.$$

6.4. $Q = E_8$**.** We give the columns of the Hilbert series matrix $H_A(t)$ of A which can be calculated from 2.4.1:

$$
\begin{aligned}
(H_A(t)_{i,1})_{1\leq i\leq 8} =& \\
= & \begin{pmatrix}
1 + t^{10} + t^{18} + t^{28} \\
t + t^9 + t^{11} + t^{17} + t^{19} + t^{27} \\
t^2 + t^8 + t^{10} + t^{12} + t^{16} + t^{18} + t^{20} + t^{26} \\
t^3 + t^7 + t^9 + t^{11} + t^{13} + t^{15} + t^{17} + t^{19} + t^{21} + t^{25} \\
t^4 + t^6 + t^8 + t^{10} + t^{12} + 2t^{14} + t^{16} + t^{18} + t^{20} + t^{22} + t^{24} \\
t^5 + t^7 + t^{11} + t^{13} + t^{15} + t^{17} + t^{21} + t^{23} \\
t^6 + t^{12} + t^{16} + t^{22} \\
t^5 + t^9 + t^{13} + t^{15} + t^{19} + t^{23}
\end{pmatrix}
\end{aligned}
$$

$$
\begin{aligned}
(H_A(t)_{i,2})_{1\leq i\leq 8} =& \\
= & \begin{pmatrix}
t + t^9 + t^{11} + t^{17} + t^{19} + t^{27} \\
1 + t^2 + t^8 + 2t^{10} + t^{12} + t^{16} + 2t^{18} + t^{20} + t^{26} + t^{28} \\
t + t^3 + t^7 + 2t^9 + 2t^{11} + t^{13} + t^{15} + 2t^{17} + 2t^{19} + t^{21} + t^{25} + t^{27} \\
t^2 + t^4 + t^6 + 2t^8 + 2t^{10} + 2t^{12} + 2t^{14} + 2t^{16} + 2t^{18} + 2t^{20} + t^{22} + t^{24} + t^{26} \\
t^3 + 2t^5 + 2t^7 + 2t^9 + 2t^{11} + 3t^{13} + 3t^{15} + 2t^{17} + 2t^{21} + 2t^{23} + t^{25} \\
t^4 + 2t^6 + t^8 + t^{10} + 2t^{12} + 2t^{14} + 2t^{16} + t^{18} + t^{20} + 2t^{22} + t^{24} \\
t^5 + t^7 + t^{11} + t^{13} + t^{15} + t^{17} + t^{21} + t^{23} \\
t^4 + t^6 + t^8 + t^{10} + t^{12} + 2t^{14} + t^{16} + t^{18} + t^{20} + t^{22} + t^{24}
\end{pmatrix},
\end{aligned}
$$

$$
(H_A(t)_{i,3})_{1\leq i\leq 8} = \begin{pmatrix}
t^2 + t^8 + t^{10} + t^{12} + t^{16} + t^{18} + t^{20} + t^{26} \\
t + t^3 + t^7 + 2t^9 + 2t^{11} + t^{13} + t^{15} \\
\quad +2t^{17} + 2t^{19} + t^{21} + t^{25} + t^{27} \\
1 + t^2 + t^4 + t^6 + 2t^8 + 3t^{10} + 2t^{12} + 2t^{14} \\
\quad +2t^{16} + 3t^{18} + 2t^{20} + t^{22} + t^{24} + t^{26} + t^{28} \\
t + t^3 + 2t^5 + 2t^7 + 3t^9 + 3t^{11} + 3t^{13} + 3t^{15} \\
\quad +3t^{17} + 3t^{19} + 2t^{21} + 2t^{23} + t^{25} + t^{27} \\
t^2 + 2t^4 + 3t^6 + 3t^8 + 3t^{10} + 4t^{12} + 4t^{14} \\
\quad +4t^{16} + 3t^{18} + 3t^{20} + 3t^{22} + 2t^{24} + t^{26} \\
t^3 + 2t^5 + 2t^7 + 2t^9 + 2t^{11} + 3t^{13} + 3t^{15} \\
\quad +2t^{17} + 2t^{19} + 2t^{21} + 2t^{23} + t^{25} \\
t^4 + t^6 + t^8 + t^{10} + t^{12} + 2t^{14} \\
\quad +t^{16} + t^{18} + t^{20} + t^{22} + t^{24} \\
t^3 + t^5 + 2t^7 + t^9 + 2t^{11} + 2t^{13} + 2t^{15} \\
\quad +2t^{17} + t^{19} + 2t^{21} + t^{23} + t^{25}
\end{pmatrix}
$$

$$(H_A(t)_{i,4})_{1\leq i\leq 8} = \begin{pmatrix} t^3 + t^7 + t^9 + t^{11} + t^{13} + t^{15} \\ \quad +t^{17} + t^{19} + t^{21} + t^{25} \\ t^2 + t^4 + t^6 + 2t^8 + 2t^{10} + 2t^{12} + 2t^{14} \\ \quad +2t^{16} + 2t^{18} + 2t^{20} + t^{22} + t^{24} + t^{26} \\ t + t^3 + 2t^5 + 2t^7 + 3t^9 + 3t^{11} + 3t^{13} + 3t^{15} \\ \quad +3t^{17} + 3t^{19} + 2t^{21} + 2t^{23} + t^{25} + t^{27} \\ 1 + t^2 + 2t^4 + 3t^6 + 3t^8 + 4t^{10} + 4t^{12} + 4t^{14} \\ \quad +4t^{16} + 4t^{18} + 3t^{20} + 3t^{22} + 2t^{24} + t^{26} + t^{28} \\ t + 2t^3 + 3t^5 + 4t^7 + 4t^9 + 5t^{11} + 5t^{13} + 5t^{15} \\ \quad +5t^{17} + 4t^{19} + 4t^{21} + 3t^{23} + 2t^{25} + t^{27} \\ t^2 + 2t^4 + 2t^6 + 3t^8 + 3t^{10} + 3t^{12} + 4t^{14} \\ \quad +3t^{16} + 3t^{18} + 3t^{20} + 2t^{22} + 2t^{24} + t^{26} \\ t^3 + t^5 + t^7 + 2t^9 + t^{11} + 2t^{13} + 2t^{15} \\ \quad +t^{17} + 2t^{19} + t^{21} + t^{23} + t^{25} \\ t^2 + t^4 + 2t^6 + 2t^8 + 2t^{10} + 3t^{12} + 2t^{14} \\ \quad +3t^{16} + 2t^{18} + 2t^{20} + 2t^{22} + t^{24} + t^{26} \end{pmatrix}$$

$$(H_A(t)_{i,5})_{1\leq i\leq 8} = \begin{pmatrix} t^4 + t^6 + t^8 + t^{10} + t^{12} + 2t^{14} \\ \quad +t^{16} + t^{18} + t^{20} + t^{22} + t^{24} \\ t^3 + 2t^5 + 2t^7 + 2t^9 + 2t^{11} + 3t^{13} + 3t^{15} \\ \quad +2t^{17} + 2t^{19} + 2t^{21} + 2t^{23} + t^{25} \\ t^2 + 2t^4 + 3t^6 + 3t^8 + 3t^{10} + 4t^{12} + 4t^{14} \\ \quad +4t^{16} + 3t^{18} + 3t^{20} + 3t^{22} + 2t24 + t^{26} \\ t + 2t^3 + 3t^5 + 4t^7 + 4t^9 + 5t^{11} + 5t^{13} + 5t^{15} \\ \quad +5t^{17} + 4t^{19} + 4t^{21} + 3t^{23} + 2t^{25} + t^{27} \\ 1 + 2t^2 + 3t^4 + 4t^6 + 5t^8 + 6t^{10} + 6t^{12} + 6t^{14} \\ \quad +6t^{16} + 6t^{18} + 5t^{20} + 4t^{22} + 3t^{24} + 2t^{26} + t^{28} \\ t + 2t^3 + 2t^5 + 3t^7 + 4t^9 + 4t^{11} + 4t^{13} + 4t^{15} \\ \quad +4t^{17} + 4t^{19} + 3t^{21} + 2t^{23} + 2t^{25} + t^{27} \\ t^2 + t^4 + t^6 + 2t^8 + 2t^{10} + 2t^{12} + 2t^{14} \\ \quad +2t^{16} + 2t^{18} + 2t^{20} + t^{22} + t^{24} + t^{26} \\ t + t^3 + 2t^5 + 2t^7 + 3t^9 + 3t^{11} + 3t^{13} + 3t^{15} \\ \quad +3t^{17} + 3t^{19} + 2t^{21} + 2t^{23} + t^{25} + t^{27} \end{pmatrix}$$

$$(H_A(t)_{i,6})_{1\leq i\leq 8} = \begin{pmatrix} t^5+t^7+t^{11}+t^{13}+t^{15}+t^{17}+t^{21}+t^{23} \\ t^4+2t^6+t^8+t^{10}+2t^{12}+2t^{14} \\ \quad +2t^{16}+t^{18}+t^{20}+2t^{22}+t^{24} \\ t^3+2t^5+2t^7+2t^9+2t^{11}+3t^{13}+3t^{15} \\ \quad +2t^{17}+2t^{19}+2t^{21}+2t^{23}+t^{25} \\ t^2+2t^4+2t^6+3t^8+3t^{10}+3t^{12}+4t^{14}+3t^{16} \\ \quad +3t^{18}+3t^{20}+2t^{22}+2t^{24}+t^{26} \\ t+2t^3+2t^5+3t^7+4t^9+4t^{11}+4t^{13} \\ \quad +4t^{15}+4t^{17}+4t^{19}+3t^{21}+2t^{23}+2t^{25}+t^{27} \\ 1+t^2+t^4+2t^6+2t^8+3t^{10}+3t^{12}+2t^{14} \\ \quad +3t^{16}+2t^{18}+2t^{20}+2t^{22}+t^{24}+t^{26}+t^{28} \\ t+t^5+t^7+t^9+2t^{11}+t^{13} \\ \quad +t^{15}+2t^{17}+t^{19}+t^{21}+t^{23}+t^{27} \\ t^2+t^4+t^6+2t^8+2t^{10}+2t^{12}+2t^{14} \\ \quad +2t^{16}+2t^{18}+2t^{20}+t^{22}+t^{24}+t^{26} \end{pmatrix}$$

$$(H_A(t)_{i,7})_{1\leq i\leq 8} = \begin{pmatrix} t^6+t^{12}+t^{16}+t^{22} \\ t^5+t^7+t^{11}+t^{13}+t^{15}+t^{17}+t^{21}+t^{23} \\ t^4+t^6+t^8+t^{10}+t^{12}+2t^{14} \\ \quad +t^{16}+t^{18}+t^{20}+t^{22}+t^{24} \\ t^3+t^5+t^7+2t^9+t^{11}+2t^{13}+2t^{15} \\ \quad +t^{17}+2t^{19}+t^{21}+t^{23}+t^{25} \\ t^2+t^4+t^6+2t^8+2t^{10}+2t^{12}+2t^{14} \\ \quad +2t^{16}+2t^{18}+2t^{20}+t^{22}+t^{24}+t^{26} \\ t+t^5+t^7+t^9+2t^{11}+t^{13}+t^{15} \\ \quad +2t^{17}+t^{19}+t^{21}+t^{23}+t^{27} \\ 1+t^6+t^{10}+t^{12}+t^{16}+t^{18}+t^{22}+t^{28} \\ t^3+t^7+t^9+t^{11}+t^{13}+t^{15} \\ \quad +t^{17}+t^{19}+t^{21}+t^{25} \end{pmatrix}$$

$$
(H_A(t)_{i,8})_{1\le i\le 8} = \begin{pmatrix}
t^5+t^9+t^{13}+t^{15}+t^{19}+t^{23} \\
t^4+t^6+t^8+t^{10}+t^{12}+2t^{14} \\
\quad +t^{16}+t^{18}+t^{20}+t^{22}+t^{24} \\
t^3+t^5+2t^7+t^9+2t^{11}+2t^{13}+2t^{15} \\
\quad +2t^{17}+t^{19}+2t^{21}+t^{23}+t^{25} \\
t^2+t^4+2t^6+2t^8+2t^{10}+3t^{12}+2t^{14} \\
\quad +3t^{16}+2t^{18}+2t^{20}+2t^{22}+t^{24}+t^{26} \\
t+t^3+2t^5+2t^7+3t^9+3t^{11}+3t^{13}+3t^{15} \\
\quad +3t^{17}+3t^{19}+2t^{21}+2t^{23}+t^{25}+t^{27} \\
t^2+t^4+t^6+2t^8+2t^{10}+2t^{12}+2t^{14} \\
\quad +2t^{16}+2t^{18}+2t^{20}+t^{22}+t^{24}+t^{26} \\
t^3+t^7+t^9+t^{11}+t^{13}+t^{15} \\
\quad +t^{17}+t^{19}+t^{21}+t^{25} \\
1+t^4+t^6+t^8+2t^{10}+t^{12}+2t^{14} \\
\quad +t^{16}+2t^{18}+t^{20}+t^{22}+t^{24}+t^{28}
\end{pmatrix}
$$

7. $HH^0(A) = Z$

From the Hilbert series 3.0.7 we see that we have one (unique up to a constant factor) central element of degree $2m_i - 2$ for each exponent $m_i < \frac{h}{2}$. We will denote a $\deg i (< h-2)$ central element by z_i.

From 2.4.2 and from the Hilbert series we can also see that the top degree $(= \deg h - 2)$ center is spanned by one element ω_i in each e_iAe_i, such that $\nu(i) = i$.

The $\omega_i \in L[h-2]$ are already given in section 5.3, and we will find the $z_i \in U[-2]$ for each Dynkin quiver separately.

7.1. $Q = D_{n+1}$.

We define the nonzero elements

$$
\begin{aligned}
b_{i,0} &= e_i, \\
b_{i,j} &= a_i^* \dots a_{i+j-1}^* a_{i+j-1} \dots a_i \text{ (where } 1 \le j \le \min\{i-1, n-1-i\}), \\
c_{i,j} &= a_i^* \dots a_{n-2}^* (a_{n-2} a_{n-2}^*) a_{n-2} \dots a_i \text{ (where } 1 \le i \le n-2,\ 1 \le j \le i-1 \\
c_{n-1,j} &= (a_{n-2} a_{n-2}^*)^j,\ 1 \le j \le n-2 \\
c_i' &= a_i^* \dots a_{n-2}^* a_{n-1}^* a_{n-1} (a_{n-2} a_{n-2}^*)^{i-1} a_{n-2} \dots a_2,\ 1 \le i \le n-1 \\
d_0 &= e_n, \\
d_j &= (a_{n-1} a_n^* a_n a_{n-1}^*)^j \text{ for } 1 \le j < \frac{n}{2}, \\
d_0' &= e_{n+1}, \\
d_j' &= (a_n a_{n-1}^* a_{n-1} a_n^*)^j \text{ for } 1 \le j < \frac{n}{2}
\end{aligned}
$$

and extend this notation for any other j, where $b_{i,j}$, $c_{i,j}$, d_j and d_j' are zero.

The exponents m_i are $1, 3, \dots, 2n-1, n$ and $h = 2n$. From 3.0.7 we get the Hilbert series of Z, depending on the parity of n, since $r_+ = n+1$ for n odd and $r_+ = n-1$ for n even:

$$n \text{ odd: } h_Z(t) = 1 + t^4 + t^8 + \dots + t^{2n-6} + (n+1)t^{2n-2},$$
$$n \text{ even: } h_Z(t) = 1 + t^4 + t^8 \dots + t^{2n-4} + (n-1)t^{2n-2}.$$

The central elements of degree $4j < 2n-2$ are

$$z_{4j} = \sum_{i=2j+1}^{n-1-2j} b_{i,2j} + \sum_{i=0}^{2j-1} c_{n-1-i,2j-i} + d_j + d'_j$$

since the relations
$b_{2j+1,2j}a_{2j} = 0$,
$a_i b_{i,2j} = b_{i+1,2j}a_i$ for $2j+1 < i < n-1-2j$,
$a_{n-1-2j}b_{n-1-2j,2j} = c_{n-2j,1}a_{n-1-2j}$,
$a_{n-1-i}c_{n-1-i,2j-1-i} = c_{n-i,2j-i}a_{n-1-i}$ for $0 < i < 2j-1$
$a_{n-1}c_{n-1,2j} = d_j a_{n-1}$, $a_n c_{n-1,2j} = d'_j a_n$
hold.

The top degree central elements are $\omega_i = c'_i$ $(1 \leq i \leq n-1)$, and additionally $\omega_n = d_{\frac{n-1}{2}}$, $-\omega_{n+1} = d'_{\frac{n-1}{2}}$ if n is odd.

We have the multiplication laws $b_{i,2j}b_{i,2k} = b_{i,2(j+k)}$, $d_j d_k = d_{j+k}$, $d'_j d'_k = d'_{j+k}$ and $c_{i,j}c_{i,k} = c_{i,j+k+n-i-1}$ which implies that $c_{n-1-i,2j-1-i}c_{n-1-i,2k-1-i} = c_{n-1-i,2j-1+2k-1+n-(n-1-i)-1} = c_{n-1-i,2(j+k)-i}$, so for $j+k < \frac{n-1}{2}$ we get the following product:

$$z_{4j}z_{4k} = z_{4(j+k)}.$$

If n is odd and $j+k = \frac{n-1}{2}$, the multiplication becomes

$$z_{4j}z_{4k} = d_{\frac{n-1}{2}} + d'_{\frac{n-1}{2}} = \omega_n - \omega_{n+1}.$$

7.2. $Q = E_6$**.** The Coxeter number is $h = 12$, and the exponents $m_i < \frac{h}{2} = 6$ are 1, 4, 5, $r_+ = 2$. For the center, we get the following Hilbert series (from Corollary 3.0.7):

$$h_Z(t) = 1 + t^6 + t^8 + 2t^{10}.$$

From the degrees, we see that the product of any two positive degree central elements is always 0. The central elements are $z_0 = 1$, z_6, z_8, ω_3 and ω_6.

We use the notation $x_i = a_i^* a_i$ and give the central elements z_6 and z_8 explicitly:

Proposition 7.2.1. *A central element of* deg 6 *is given by*

$$z_6 = a_1 a_2 x_3 a_2^* a_1^* - a_2 x_3^2 a_2^* - x_5 x_3 x_5 + a_3 x_2^2 a_3^* - a_4 a_3 x_2 a_3^* a_4^*$$

Proof. Observe that $z_6^* = z_6$, so it is enough to show that z_6 commutes with all a_i. Since $\eta(z_6) = z_6$, we have only to show that z_6 commutes with a_1, a_2, a_5:

$$\begin{aligned} a_1 z_6 &= -a_1 a_2 x_3^2 a_2^* = -a_1 a_2 x_5 x_2 a_2^* = a_1 a_2 x_3 a_2^* a_1^* a_1 = z_6 a_1, \\ a_2 z_6 &= -a_2 x_5 x_3 x_5 = -a_2 x_3 x_2 x_5 = -a_2 x_3 x_3 x_2 = z_6 a_2, \\ a_5 z_6 &= a_5 x_5 x_3 x_5 = 0 = z_6 a_5. \end{aligned}$$

□

Proposition 7.2.2. *The* deg 8 *central element of* A *is given by*

$$z_8 = -a_2 x_5 x_3 x_5 a_2^* - x_5 x_3^2 x_5 - a_3 x_5 x_2 x_5 a_3^*.$$

Proof. Since $z_8^* = z_8$ and $\eta(z_8) = z_8$, it is enough to show that z_8 commutes with a_1, a_2 and a_5:

$$\begin{aligned} a_1 z_8 &= -a_1 a_2 x_5 x_3 x_5 a_2^* = -a_1 a_2 x_5 x_2 x_3 a_2^* = -a_1 a_2 x_2 x_3 x_3 a_2^* = 0 = a_1 z_8, \\ a_2 z_8 &= -a_2 x_5 x_3^2 x_5 = -a_2 x_5 x_2 x_3 x_2 = -a_2 x_5 x_3 x_5 x_2 = z_8 a_2, \\ a_5 z_8 &= a_5 x_5 x_3 x_5 x_3 = 0 = z_8 a_5. \end{aligned}$$

□

7.3. $Q = E_7$**.** The Coxeter number is $h = 18$, the exponents $m_i < \frac{h}{2} = 9$ are $1, 5, 7$, $r_+ = 7$, and the Hilbert series of the center is (see Corollary 3.0.7):

$$h_Z(t) = 1 + t^8 + t^{12} + 7t^{16}$$

The center is spanned by $z_0 = 1$, z_8, z_{12}, $\omega_1, \ldots, \omega_7$. The only interesting product to compute is z_8^2 which lies in the top degree.

We give z_8 and z_{12} explicitly:

Proposition 7.3.1. *The* deg 8 *central element of* A *is given by*

$$\begin{aligned} z_8 = &-a_1 a_2 a_3 x_6 a_3^* a_2^* a_1^* - a_2 a_3 x_4^2 a_3^* a_2^* - a_3 x_6 x_4 x_6 a_3^* - x_4 x_3^2 x_4 \\ &- a_4 x_4 x_6 x_4 a_4^* + a_6 x_4 x_6 x_4 a_6^*. \end{aligned}$$

Proof. It is clear that $z_8^* = z_8$, so it is enough to prove that z_8 commutes with every a_i:

$$\begin{aligned} a_1 z_8 &= -a_1 a_2 a_3 x_4^2 a_3^* a_2^* = -a_1 a_2 a_3 x_6 x_3 a_3^* a_2^* = -a_1 a_2 a_3 x_6 a_3^* a_2^* x_1 = z_8 a_1, \\ a_2 z_8 &= -a_2 a_3 x_6 x_4 x_6 a_3^* = -a_2 a_3 x_4 x_3 x_6 a_3^* = -a_2 a_3 x_4^2 x_3 a_3^* = -a_2 a_3 x_4^2 a_3^* x_2 = z_8 a_2, \\ a_3 z_8 &= -a_3 x_4 x_3^2 x_4 = -a_3 x_6 x_4 x_3 x_4 = -a_3 x_6 x_4 x_6 x_3 = z_8 a_3, \\ a_4 z_8 &= -a_4 x_4 x_3^2 x_4 = -a_4 x_4 x_6 x_4^2 = z_8 a_4 \\ a_5 z_8 &= -a_5 a_4 x_4 x_6 x_4 a_4^* = 0 = z_8 a_5 \\ a_6 z_8 &= -a_6 x_4 x_3^2 x_4 = -a_6 x_3^3 x_6 = a_6 x_4 x_6 x_4 x_6 = z_8 a_6. \end{aligned}$$

□

Proposition 7.3.2. *The* deg 12 *central element of* A *is*

$$z_{12} = -a_3x_4x_6x_4x_6x_4a_3^* - x_4x_6x_4^2x_6x_4 + a_4x_6x_4x_6x_4x_6a_4^* + a_6x_4x_6x_4x_6x_4a_6^*.$$

Proof. $z_{12}^* = z_{12}$ is clear, so we have only to show that z_{12} commutes with all a_i:

$$\begin{aligned}
a_1z_{12} &= 0 = z_{12}a_1,\\
a_2z_{12} &= -a_2a_3x_4x_6x_4x_6x_4a_3^* = a_2a_3x_4x_6x_3^2x_6a_3^* = a_2a_3x_4x_6x_4^2x_6a_3^*\\
&= a_2a_3x_3^2x_4^2x_6a_3^* = 0 = z_{12}a_2,\\
a_3z_{12} &= -a_3x_4x_6x_4^2x_6x_4 = a_3x_4x_6x_3^2x_6x_3 = -a_3(x_4x_6)^2x_4x_3 = z_{12}a_3,\\
a_4z_{12} &= -a_4x_4x_6x_4^2x_6x_4 = a_4x_3^2x_4x_3x_6x_4 = a_4(x_6x_4)^2x_6 = z_{12}a_4,\\
a_5z_{12} &= a_5a_4x_6x_4x_6x_4x_6a_4^* = -a_5a_4x_6x_3^3x_6a_4^* = a_5a_4x_3^4x_6a_4^* = 0 = z_{12}a_5,\\
a_6z_{12} &= -a_6x_4x_6x_4^2x_6x_4 = a_6x_4x_6x_3^2x_6x_3 = -a_6x_4x_6x_4x_6x_4x_3 =\\
&= a_6(x_4x_6)^3 + a_6(x_4x_6)^2x_4^2 = z_{12}a_6 + a_6x_3^4x_4^2 = z_{12}a_6.
\end{aligned}$$

□

Now we calculate the product z_8^2:

Proposition 7.3.3.

$$z_8^2 = \omega_1 + \omega_3 - \omega_7.$$

Proof. We get

$$\begin{aligned}
z_8^2 &= (a_1a_2a_3x_6a_3^*a_2^*a_1^*)^2 + (a_2a_3x_4^2a_3^*a_2^*)^2 + (a_3x_6x_4x_6a_3^*)^2 + (x_4x_3^2x_4)^2\\
&\quad + (a_4x_4x_6x_4a_4^*)^2 + (a_6x_4x_6x_4a_6^*)^2\\
&= \underbrace{a_1a_2a_3x_6x_3^3x_6a_3^*a_2^*a_1^*}_{=\omega_1} + \underbrace{a_2a_3x_4^2x_3^2x_4^2a_3^*a_2^*}_{=a_2a_3x_4^2x_6^2x_4^2a_3^*a_2^*=0} + \underbrace{a_3x_6x_4x_6x_3x_6x_4x_6a_3^*}_{=\omega_3}\\
&\quad + \underbrace{x_4x_3^2x_4^3x_3^2x_4}_{=0} + \underbrace{a_4x_4x_6x_4^3x_6x_4a_4^*}_{=0} + \underbrace{a_6(x_4x_6)^3x_4a_6^*}_{=-\omega_7}
\end{aligned}$$

□

7.4. $Q = E_8$**.** The Coxeter number $h = 30$, and the exponents $m_i < \frac{h}{2} = 15$ are $1, 7, 11, 13$, $r_+ = 8$. For the center, we get the following Hilbert series (from Corollary 3.0.7):

$$h_Z(t) = 1 + t^{12} + t^{20} + t^{24} + 8t^{28}.$$

The center is spanned by $z_0 = 1$, z_{12}, z_{20}, z_{24}, $\omega_1, \ldots, \omega_8$. The only interesting product is z_{12}^2.

Proposition 7.4.1. *The* deg 12 *central element of* A *is*

$$\begin{aligned}
z_{12} &= a_1a_2a_3x_6x_4x_6a_3^*a_2^*a_1^* + a_2a_3x_4x_3^2x_4a_3^*a_2^* + a_3(x_4x_6)^2x_4a_3^*\\
&\quad + (x_3x_4x_3)^2 - a_4(x_6x_4)^2x_6a_4^* + a_5a_4x_6x_4^2x_6a_4^*a_5^* - a_6(x_4x_6)^2x_4a_6^*.
\end{aligned}$$

Proof. $z_{12}^* = z_{12}$, so we only have to show that z_{12} commutes with all a_i:

$$\begin{aligned}
a_0 z_{12} &= a_0 a_1 a_2 a_3 x_6 x_4 x_6 a_3^* a_2^* a_1^* = a_0 a_1 a_2 a_3 x_6 x_3 x_4 a_3^* a_2^* a_1^* \\
&= a_0 a_1 a_2 a_3 x_3 x_4^2 a_3^* a_2^* a_1^* = 0 = z_{12} a_0, \\
a_1 z_{12} &= a_1 a_2 a_3 x_4 x_3^2 x_4 a_3^* a_2^* = a_1 a_2 a_3 x_6 x_4 x_6 x_3 a_3^* a_2^* \\
&= a_1 a_2 a_3 x_6 x_4 x_6 a_3^* a_2^* x_1 = z_{12} a_1, \\
a_2 z_{12} &= a_2 a_3 (x_4 x_6)^2 x_4 a_3^* = -a_2 a_3 x_4 x_3^2 x_6 x_3 a_3^* = a_2 a_3 x_4 x_3^2 x_4 a_3^* x_2 = z_{12} a_2, \\
a_3 z_{12} &= a_3 (x_3 x_4 x_3)^2 = -a_3 x_3 x_6 x_3^2 x_4 x_3 = a_3 x_4 x_6 x_4 x_6 x_4 x_3 = z_{12} a_3, \\
a_4 z_{12} &= a_4 (x_3 x_4 x_3)^2 = -a_4 (x_6 x_4)^3 = z_{12} a_4, \\
a_5 z_{12} &= -a_5 a_4 (x_6 x_4)^2 x_6 a_4^* = a_5 a_4 x_3^4 x_6 a_4^* = -a_5 a_4 x_3^4 x_4 a_4^* \\
&= a_5 a_4 x_6 x_4^2 x_6 a_4^* x_5 = z_{12} a_5, \\
a_6 z_{12} &= a_6 (x_3 x_4 x_3)^2 = -a_6 (x_6 x_4)^2 x_6 = z_{12} a_6.
\end{aligned}$$

□

Proposition 7.4.2. *The central element of* deg 20 *is*

$$\begin{aligned}
z_{20} = &-a_1 a_2 a_3 x_4^2 x_3^3 x_4^2 a_3^* a_2^* a_1^* - a_2 a_3 (x_6 x_4)^2 (x_4 x_6)^2 a_3^* a_2^* + a_3 (x_6 x_4)^4 x_6 a_3^* \\
&- (x_4 x_6)^5 + (x_6 x_4^2)^3 x_6 - (x_6 x_4)^5 \\
&- a_4 (x_4 x_6 x_4)^3 a_4^* - a_6 (x_4 x_6)^4 x_4 a_6^*.
\end{aligned}$$

Proof. $z_{20}^* = z_{20}$, so it is sufficient to show that z_{20} commutes with all a_i:

$$\begin{aligned}
a_0 z_{20} &= -a_0 a_1 a_2 a_3 x_4^2 x_3^3 x_4^2 a_3^* a_2^* a_1^* = a_0 a_1 a_2 a_3 x_4^2 x_6 x_4 x_6 x_4^2 a_3^* a_2^* a_1^* \\
&= -a_0 a_1 a_2 a_3 x_4^2 x_6 x_4 x_6 x_4 x_6 a_3^* a_2^* a_1^* = -a_0 a_1 a_2 a_3 x_4^2 x_3^4 x_6 a_3^* a_2^* a_1^* \\
&= -a_0 a_1 a_2 a_3 x_6 x_3^5 x_6 a_3^* a_2^* a_1^* = 0 = z_{20} a_0, \\
a_1 z_{20} &= -a_1 a_2 a_3 (x_6 x_4)^2 (x_4 x_6)^2 a_3^* a_2^* = -a_1 a_2 a_3 x_4 x_3 x_6 x_4 (x_4 x_6)^2 a_3^* a_2^* \\
&= -a_1 a_2 a_3 x_4^2 x_3^3 x_4 x_3 x_6 a_3^* a_2^* = -a_1 a_2 a_3 x_4^2 x_3^3 x_4^2 x_3 a_3^* a_2^* \\
&= -a_1 a_2 a_3 x_4^2 x_3^3 x_4^2 a_3^* a_2^* x_1 = z_{20} a_1, \\
a_2 z_{20} &= a_2 a_3 (x_6 x_4)^4 x_6 a_3^* = a_2 a_3 x_6 x_4 x_6 x_3^3 x_6 x_3 x_6 a_3^* \\
&= -a_2 a_3 x_6 x_4 x_6 x_3^3 x_4^2 x_3 a_3^* = a_2 a_3 (x_6 x_4)^2 x_4 x_6 x_4^2 x_3 a_3^* \\
&= -a_2 a_3 (x_6 x_4)^2 (x_4 x_6)^2 a_3^* x_2 = z_{20} a_2.
\end{aligned}$$

From

$$a_3 (x_6 x_4^2)^3 x_6 = a_3 x_4 x_3^3 (x_4^2 x_6)^2 = a_3 x_4 x_6 x_4 x_6 x_3^2 x_4 x_3^2 x_6 = a_3 (x_4 x_6)^5$$

follows

$$a_3 z_{20} = a_3 (-(x_4 x_6)^5 + (x_6 x_4^2)^3 x_6 - (x_6 x_4)^5) = a_3 (x_6 x_4)^4 x_6 x_3 = z_{20} a_3.$$

To show that z_{20} commutes with a_4, we use the following rules:

$$\begin{aligned} a_4(x_4x_6)^3 &= a_4x_4x_3^4x_6 \\ &= -a_4x_6x_4^2x_6x_4x_6 - a_4x_6x_4x_6x_4^2x_6 \\ &= -a_4(x_4x_6x_4)^2 - a_4(x_4x_6)^2x_4^2 \qquad = a_4(x_6x_4^2)^2 + a_4(x_6x_4)^2, \end{aligned}$$

so

$$\begin{aligned} a_4(x_4x_6)^5 &= -a_4x_6x_4^2x_6(x_4x_6)^3 - a_4(x_6x_4x_6x_4)^2(x_4x_6)^3 \\ &= a_4x_6x_4(x_4x_6x_4)^2x_4x_6 + a_4x_6x_4(x_4x_6)^2x_4^3x_6 \\ &\quad - a_4(x_6x_4)^2(x_6x_4^2)^2 - a_4(x_6x_4)^5, \end{aligned}$$

$$\begin{aligned} a_4z_{20} &= -a_4(x_4x_6)^5 + a_4(x_6x_4^2)^3x_6 - a_4(x_6x_4)^5 = a_4(x_6x_4)^2(x_6x_4^2)^2 \\ &= a_4x_6x_3^4x_4^2x_6x_4^2 = a_4(x_4x_6x_4)^3x_4 = z_{20}a_4, \\ a_5z_{20} &= -a_5a_4(x_4x_6x_4)^3a_4^* = 0 = z_{20}a_5, \\ a_6z_{20} &= -a_6(x_4x_6)^5 + a_6(x_6x_4^2)^3x_6 - a_6(x_6x_4)^5 = -a_6(x_4x_6)^5 = z_{20}a_6. \end{aligned}$$

□

In deg 24 there is exactly one central element (up to constant multiple). Since we already have a deg 12 central element, the most obvious candidate is its square. In fact, this is true from the following

Proposition 7.4.3. *The* deg 24 *central element of* A *is*

$$z_{24} = z_{12}^2.$$

Proof. A simple calculation shows that z_{12}^2 is nonzero:

$$x_4x_3z_{12}^2 = \omega_4.$$

□

8. $HH^1(A)$

Recall Theorem 3.0.6 where we know that $HH^1(A)$ is isomorphic to the non-topdegree part of $HH^0(A)$. In fact, $HH^1(A)$ is generated by the central elements in the following way:

Proposition 8.0.4. $HH^1(A)$ *is spanned by maps*

$$\begin{gathered} \theta_k : (A \otimes V \otimes A) \longrightarrow A, \\ \theta_k(1 \otimes a_i \otimes 1) = 0, \\ \theta_k(1 \otimes a_i^* \otimes 1) = a_i^* z_k. \end{gathered}$$

Proof. These maps clearly lie in $\ker d_2^*$: Recall

$$\begin{array}{ccc} A \otimes A & \xrightarrow{d_2} & A \otimes V \otimes A \\ x \otimes y & \longmapsto & \sum\limits_{a \in \bar{Q}} \epsilon_a xa \otimes a^* \otimes y + \sum\limits_{a \in \bar{Q}} \epsilon_a x \otimes a \otimes a^* y, \end{array}$$

then

$$\begin{aligned} d_2^* \circ \theta_k(1 \otimes 1) &= \theta_k(\sum_{a \in \bar{Q}} \epsilon_a a \otimes a^* \otimes 1 + \sum_{a \in \bar{Q}} \epsilon_a 1 \otimes a \otimes a^*) \\ &= \sum_{i \in I} a_i a_i^* z_k - \sum_{i \in I} a_i^* a_i z_k = \sum_{i \in I} [a_i, a_i^*] z_k = 0. \end{aligned}$$

We will later see in section 11 that $HH^4(A)$ is generated by ζ_k where $\zeta_k(\theta_k) = 1$ under the duality $HH^4(A) = (HH^1(A))^*$ established in [EE2], so θ_k is nonzero in $HH^1(A)$. □

9. $HH^2(A)$

We know from 3.0.6 that $HH^2(A) = K[-2]$ lies in degree -2, i.e. in the lowest degree of $A^R[-2]$ (using the identifications in [EE2, Section 4.5.]), that is in $R[-2]$. Since the image of d_2^* lies in degree > -2, $HH^2(A) = \ker d_3^*$.

Proposition 9.0.5. *$HH^2(A)$ is given by the kernel of the matrix $H_A(1)$, where we identify $\mathbb{C}^I = R = \oplus_{i \in I} Re_i$.*

Proof. Recall

$$d_3^*(x) = \sum_{x_i \in B} x_i x x_i^* = \sum_{j,k \in I} \sum_{x_i \in B_{j,k}} x_i x x_i^*.$$

For each $x_i \in e_k A e_j$, we see that $x_i e_l x_i^* = \delta_{jl} \omega_k$.

It follows that for $x = \sum\limits_{i \in I} \lambda_i e_i$ the map is given by

$$d_3^*(x) = \sum_{i \in I} \mu_i \omega_i,$$

where the vectors $\lambda = (\lambda_i)_{i \in I} \in \mathbb{C}^I$ and $\mu = (\mu_i)_{i \in I} \in \mathbb{C}^I$ satisfy the equation

$$H_A(1)\lambda = \mu. \tag{9.0.6}$$

So the kernel of d_3^* is given by the kernel of $H_A(1)$. □

Now, we find the elements in $HH^2(A)$ for the quivers separately.

9.1. $Q = D_{n+1}$, n even.

$$H_A(1) = \begin{pmatrix} 2 & 2 & 2 & 2 & \dots & \dots & 2 & 1 & 1 \\ 2 & 4 & 4 & 4 & \dots & \dots & 4 & 2 & 2 \\ 2 & 4 & 6 & 6 & \dots & \dots & 6 & 3 & 3 \\ 2 & 4 & 6 & 8 & \dots & \dots & 8 & 4 & 4 \\ \vdots & \vdots & \vdots & \vdots & \ddots & & \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \vdots & & \ddots & \vdots & \vdots & \vdots \\ 2 & 4 & 6 & 8 & \dots & \dots & 2(n-1) & n-1 & n-1 \\ 1 & 2 & 3 & 4 & \dots & \dots & n-1 & \frac{n}{2} & \frac{n}{2} \\ 1 & 2 & 3 & 4 & \dots & \dots & n-1 & \frac{n}{2} & \frac{n}{2} \end{pmatrix} \tag{9.1.1}$$

with kernel $\langle e_n - e_{n+1}\rangle$. So a basis of $HH^2(A)$ is given by

$$\{f_n = [e_n - e_{n+1}]\}.$$

9.2. $Q = E_6$.

$$\begin{pmatrix} 2 & 3 & 4 & 3 & 2 & 2 \\ 3 & 6 & 8 & 6 & 3 & 4 \\ 4 & 8 & 12 & 8 & 4 & 6 \\ 3 & 6 & 8 & 6 & 3 & 4 \\ 2 & 3 & 4 & 3 & 2 & 2 \\ 2 & 4 & 6 & 4 & 2 & 4 \end{pmatrix} \tag{9.2.1}$$

with kernel $\langle e_1 - e_5, e_2 - e_4\rangle$. So a basis of $HH^2(A)$ is given by

$$\{f_1 = [e_1 - e_5], f_2 = [e_2 - e_4]\}.$$

10. $HH^3(A)$

We know that $HH^3(A)$ lives in degree -2. Using the notations and identifications in [EE2, Section 4.5.], we see that the kernel of d_4^* has to be the top degree part of $\mathcal{N}^R[-h]$ (since $Im\, d_3^*$ lives in degree -2), so

$$HH^3(A) = \mathcal{N}^R[-h](-2)/Im\, d_3^*.$$

Proposition 10.0.2. *$HH^3(A)$ is given by the cokernel of the matrix $H_A(1)$, where we identify $\mathbb{C}^I = A^{top} = \oplus_{i\in I} e_i A^{top} e_{\nu(i)}$.*

Proof. This follows immediately from the discussion in the previous section because d_3^* is given by $H_A(1)$. □

Note that $HH^3(A) = (HH^2(A))^*$ under the duality in [EE2]. We choose a basis h_i of $HH^3(A)$, so that $h_i(f_j) = \delta_{i,j}$

10.1. $Q = D_{n+1}$, n even. From $H_A(1)$ in 9.1.1 we see that:

$$\begin{array}{rcll} d_3^*(2e_1 - e_2) & = & 2\omega_1, & \\ d_3^*(-e_{i-1} + 2e_i - e_{i+1}) & = & 2\omega_i & \forall 2 \le i \le n-2, \\ d_3^*((-n-1)e_{n-2} + 2(n-1)e_{n-1} - 2(n-1)e_n) & = & (n-1)\omega_{n-1}, & \\ d_3^*(2e_n - e_{n-1}) & = & \omega_n + \omega_{n+1}, & \end{array}$$

so

$$HH^3(A) = (\mathcal{N}^R)^{top}[-h]/(\omega_1 = \omega_2 = \ldots = \omega_{n-1} = 0,\ \omega_n + \omega_{n+1} = 0)$$

with basis

$$\{h_n = [\omega_n]\}$$

10.2. $Q = E_6$**.** From $H_A(1)$ in 9.2.1 we see that:

$$\begin{aligned} d_3^*(2e_1 - e_2) &= \omega_1 + \omega_5, \\ d_3^*(-e_1 + 2e_2 - e_3) &= \omega_2 + \omega_4, \\ d_3^*(-2e_2 + 2e_3 - e_6) &= 2\omega_3, \\ d_3^*(-e_3 + 2e_6) &= 2\omega_6, \end{aligned}$$

so

$$HH^3(A) = (\mathcal{N}^R)^{top}[-h]/(\omega_3 = \omega_6 = \omega_1 + \omega_5 = \omega_2 + \omega_3 = 0)$$

with basis

$$\{h_1 = [\omega_1],\ h_2 = [\omega_2]\}.$$

11. $HH^4(A)$

We have $HH^4(A) = U^*[-2]$, so its top degree is -4, and its generators sit in degrees $-4 - \deg z_k$ for each central element, one in each degree.

Proposition 11.0.1. *Let* $\zeta_0 \in \ker d_5^*$ *be a top degree element in* $(V \otimes \mathcal{N})^R[-h-2]$*, such that* $m(\zeta_0)$ *is nonzero, where* m *is the multiplication map. Then* $HH^4(A)$ *is generated by elements* $\zeta_k \in \ker d_5^*$ *which satisfy* $\zeta_k z_k = \zeta_0$*.*

Proof. If $x \in \mathcal{N}^R[-h])$ lies in degree -4, then $m(d_4^*(x)) = 0$, so ζ_0 is nonzero in $HH^4(A)$.

For every non-topdegree central element z_k we can find a ζ_k satisfying the properties above, which is done for each quiver separately below.

For any central element $z \in A$, we have that $d_4^*(zy) = d_4^*(y)z$. If $\zeta_k = d_4^*(y)$, then by construction $\zeta_0 = \zeta_k z_k = d_4^*(z_k y)$ which is a contradiction.

So these ζ_k are all nonzero in $HH^4(A)$, and also generate this cohomology space.

□

A basis of $HH^4(A)$ is given by these ζ_k, and we choose them so that $\zeta_k(\theta_k) = 1$ under the duality $HH^4(A) = (HH^1(A))^*$ in [EE2].

11.1. $Q = D_{n+1}$**,** n **odd.** We define

$$\begin{aligned} \zeta_0 =& [a_{n-1}^* \otimes a_{n-1}a_n^*a_n(a_{n-1}^*a_{n-1}a_n^*a_n)^{\frac{n-3}{2}} + \quad a_{n-1} \otimes a_n^*a_na_{n-1}^*(a_{n-1}a_n^*a_na_{n-1}^*)^{\frac{n-3}{2}}], \\ \zeta_{4k} =& \frac{1}{2}[a_{n-1}^* \otimes a_{n-1}a_n^*a_n(a_{n-1}^*a_{n-1}a_n^*a_n)^{\frac{n-3}{2}-k} + \quad a_{n-1} \otimes a_n^*a_na_{n-1}^*(a_{n-1}a_n^*a_na_{n-1}^*)^{\frac{n-3}{2}-k} - \\ & a_n^* \otimes a_na_{n-1}^*a_{n-1}(a_n^*a_na_{n-1}^*a_{n-1})^{\frac{n-3}{2}-k} - \quad a_n \otimes a_{n-1}^*a_{n-1}a_n^*(a_na_{n-1}^*a_{n-1}a_n^*)^{\frac{n-3}{2}-k}]. \end{aligned}$$

11.2. $Q = D_{n+1}$**,** n **even.** We define

$$\begin{aligned}
\zeta_0 =&[a_{n-1}^* \otimes a_{n-1}(a_n^* a_n a_{n-1}^* a_{n-1})^{\frac{n-2}{2}-k} \qquad + a_{n-1} \otimes a_n^*(a_n a_{n-1}^* a_{n-1} a_n^*)^{\frac{n-2}{2}-k}],\\
\zeta_{4k} =&\frac{1}{2}[a_{n-1}^* \otimes a_{n-1}(a_n^* a_n a_{n-1}^* a_{n-1})^{\frac{n-2}{2}-k} \quad + a_{n-1} \otimes a_n^*(a_n a_{n-1}^* a_{n-1} a_n^*)^{\frac{n-2}{2}-k} -\\
& a_n^* \otimes a_n(a_{n-1}^* a_{n-1} a_n^* a_n)^{\frac{n-2}{2}-k} - \qquad a_n \otimes a_{n-1}^*(a_{n-1} a_n^* a_n a_{n-1}^*)^{\frac{n-2}{2}-k}].
\end{aligned}$$

11.3. $Q = E_6$**.** We define

$$\begin{aligned}
\zeta_0 &= [a_3^* \otimes a_3(a_2^* a_2 a_3^* a_3)^2 + a_3 \otimes a_2^*(a_2 a_3^* a_3 a_2^*)^2],\\
\zeta_6 &= \frac{1}{4}[-a_3^* \otimes a_3 a_2^* a_2 - a_3 \otimes a_2^* a_2 a_2^* + a_2^* \otimes a_2 a_2^* a_2 + a_2 \otimes a_2^* a_2 a_3^*\\
&\qquad -a_2^* \otimes a_2 a_3^* a_3 - a_2 \otimes a_3^* a_3 a_3^* + a_3^* \otimes a_3 a_3^* a_3 + a_3 \otimes a_3^* a_3 a_2^*],\\
\zeta_8 &= \frac{1}{2}[a_3^* \otimes a_3 + a_3 \otimes a_2^* - a_2^* \otimes a_2 - a_2 \otimes a_3^*].
\end{aligned}$$

11.4. $Q = E_7$**.** We define

$$\begin{aligned}
\zeta_0 &= [a_4^* \otimes a_4 a_3^* a_3(a_4^* a_4 a_3^* a_3)^3 + a_4 \otimes a_3^* a_3 a_4^*(a_4 a_3^* a_3 a_4^*)^3],\\
\zeta_8 &= \frac{1}{2}[a_4^* \otimes a_4 a_3^* a_3 a_4^* a_4 a_3^* a_3 + a_4 \otimes a_3^* a_3 a_4^* a_4 a_3^* a_3 a_4^*\\
&\quad -a_3^* \otimes a_3 a_4^* a_4 a_3^* a_3 a_4^* a_4 - a_3 \otimes a_4^* a_4 a_3^* a_3 a_4^* a_4 a_3^*],\\
\zeta_{12} &= \frac{1}{2}[a_4^* \otimes a_4 a_3^* a_3 + a_4 \otimes a_3^* a_3 a_4^* - a_3^* \otimes a_3 a_4^* a_4 - a_3 \otimes a_4^* a_4 a_3^*].
\end{aligned}$$

11.5. $Q = E_8$**.** We define

$$\begin{aligned}
\zeta_0 &= [a_4^* \otimes a_4 a_3^* a_3(a_4^* a_4 a_3^* a_3)^6 + a_4 \otimes a_3^* a_3 a_4^*(a_4 a_3^* a_3 a_4^*)^6],\\
\zeta_{12} &= \frac{1}{2}[a_4^* \otimes a_4 a_3^* a_3(a_4^* a_4 a_3^* a_3)^3 + a_4 \otimes a_3^* a_3 a_4^*(a_4 a_3^* a_3 a_4^*)^3\\
&\quad -a_3^* \otimes a_3 a_4^* a_4(a_3^* a_3 a_4^* a_4)^3 - a_3 \otimes a_4^* a_4 a_3^*(a_3 a_4^* a_4 a_3^*)^3],\\
\zeta_{20} &= \frac{1}{2}[a_4^* \otimes a_4 a_3^* a_3 a_4^* a_4 a_3^* a_3 + a_4 \otimes a_3^* a_3 a_4^* a_4 a_3^* a_3 a_4^*\\
&\quad -a_3^* \otimes a_3 a_4^* a_4 a_3^* a_3 a_4^* a_4 - a_3 \otimes a_4^* a_4 a_3^* a_3 a_4^* a_4 a_3^*],\\
\zeta_{24} &= \frac{1}{2}[a_4^* \otimes a_4 a_3^* a_3 + a_4 \otimes a_3^* a_3 a_4^* - a_3^* \otimes a_3 a_4^* a_4 - a_3 \otimes a_4^* a_4 a_3^*].
\end{aligned}$$

12. $HH^5(A)$

We have $HH^5(A) = U^*[-2] \oplus Y^*[-h-2]$. We discuss these two subspaces separately.

12.1. $U^*[-2]$**.** In $U^*[-2]$, like in $HH^4(A)$, we have generators coming from the center in some dual sense.

We have $d_6^*(U^*[-2]) = 0$.

Proposition 12.1.1. *Let* ψ_0 *be a top degree element* $[\omega_i]$ *in some* $e_i \mathcal{N}^R e_i[-h-2]$*. Then* $HH^5(A)$ *is generated by* $\psi_k \in \mathcal{N}^R$ *which satisfy* $\psi_k z_k = \psi_0$*.*

Proof. If $\sum_{a\in\bar{Q}} a\otimes x_a \in V\otimes\mathcal{N}^R$ lies in degree -4, then the image of $d_5^*(x) = \sum_a ax_a - x_a\eta(a)$, under the linear map f (which is associated to A as a Frobenius algebra) is zero where $f(\omega_i) = 1$. So ψ_0 is nonzero in $HH^5(A)$.

For every non-topdegree central element z_k we can find a ζ_k satisfying the properties above, which is done for each quiver separately in subsection 12.3.

For any central element $z \in A$, we have that $d_5^*(zy) = d_5^*(y)z$. If $\psi_k = d_5^*(y)$, then by construction $\psi_0 = \psi_k z_k = d_4^*(z_k y)$ which is a contradiction.

So these ψ_k are nonzero in $HH^5(A)$ and generate this cohomology space. □

The relation $ax_a = x_a\eta(a)$ then gives us that all ω_i's are equivalent in $HH^5(A)$.

12.2. $Y^*[-h-2]$.

We have to introduce some new notations.

Definition 12.2.1. *We define F to be the set of vertices in I which are fixed by $\nu(i)$, i.e.*

$$F = \{i \in I|\nu(i) = i\}.$$

Definition 12.2.2. *Let η_{ij} be the restriction of η on e_iAe_j $(i,j\in F)$. Let $n_{ij}^+ = \dim\ker(\eta_{ij}-1)$ and $n_{ij}^- = \dim\ker(\eta_{ij}+1)$.*

We define the signed truncated dimension matrix $(H_A^\eta)_{i,j\in F}$ *in the following way:*

$$(H_A^\eta)_{ij} = n_{ij}^+ - n_{ij}^-.$$

Now we can make the following statement:

Proposition 12.2.3. *$Y^*[-h-2]$ is given by the kernel of the matrix H_A^η, where we identify $\mathbb{C}^F = \oplus_{i\in F} Re_i$.*

Proof. $Y^*[-h-2]$ is the kernel of the restriction $d_6^*|_{\mathcal{N}^R[-h-2](-h-2)=R_F[-h-2]} \to A^R[-2h]$, where R_F is the linear span of e_i's, such that i is fixed by ν,

$$d_6^*(x) = \sum_{x_j\in B} x_j x\eta(x_j^*) = \sum_{x_j\in B}\eta(x_j)xx_j^*.$$

then

$$d_6^* : R_F[-h-2] \to (A^{top})^R[-2h]$$

can also be written as a matrix multiplication

$$H_A^\eta : \mathbb{C}^F \to \mathbb{C}^F$$

under the identifications $R_F = \mathbb{C}^F = \bigoplus_{i\in F} e_iA^{top}e_i$. □

We compute the matrices H_A^η and their kernels for each quiver separately.

Recall that $\dim Y = r_+ - r_- - \#\{m_i|m_i = \frac{h}{2}\} = \dim R_F - \#\{m_i|m_i = \frac{h}{2}\}$. We will find Y^* explicitely for each quiver.

12.2.1. $Q = E_6, E_8$. $\frac{h}{2}$ is not an exponent, so $Y^* = R_F$.

12.2.2. $Q = D_{n+1}$, *n odd.* All basis elements of e_kAe_j given in section 6 are eigenvectors of η_{kj}.

For any of these basis elements x, $\eta(x) = (-1)^{n_x}x$ where n_x is the number of no-star letters in the monomial expression of x. So H_A^η can be computed directly.

For $k \leq j \leq n-1$:

$$\begin{aligned}(H_A^\eta)_{k,j} &= \begin{cases} \sum\limits_{l=0}^{k-1}(-1)^l + (-1)^{n-j}\sum\limits_{l=0}^{k-1}(-1)^l & k \leq n-j \\ \begin{array}{c}\sum\limits_{l=0}^{n-j-1}(-1)^l + (-1)^{n-j}\sum\limits_{l=0}^{k-1}(-1)^l \\ +(-1)^{n-j}\sum\limits_{l=0}^{k-1+j-n}(-1)^l\end{array} & k > n-j \end{cases} \\ &= \begin{cases} 2 & k, l \; odd \\ 0 & else \end{cases}.\end{aligned}$$

For $j < k \leq n-1$:

$$\begin{aligned}(H_A^\eta)_{k,j} &= \begin{cases} (-1)^{k-j}\sum\limits_{l=0}^{j-1}(-1)^l + (-1)^{n-j}\sum\limits_{l=0}^{j-1}(-1)^l & j \leq n-k \\ \begin{array}{c}(-1)^{k-j}\sum\limits_{l=0}^{n-k-1}(-1)^l + (-1)^{n-j}\sum\limits_{l=0}^{j-1}(-1)^l \\ +(-1)^{n-j}\sum\limits_{l=0}^{k-1+j-n}(-1)^l\end{array} & j > n-k \end{cases} \\ &= \begin{cases} 2 & k, l \; odd \\ 0 & else \end{cases}.\end{aligned}$$

For $k, j \leq n-1$:

$$\begin{aligned}(H_A^\eta)_{k,n} &= \sum_{l=0}^{k-1}(-1)^l = \begin{cases} 1 & k \; odd \\ 0 & k \; even \end{cases}, \\ (H_A^\eta)_{n,n} &= \sum_{l=0}^{\frac{n-1}{2}}(-1)^{2l} = \frac{n+1}{2}, \\ (H_A^\eta)_{n+1,n} &= \sum_{l=0}^{\frac{n-3}{2}}(-1)^{1+2l} = -\frac{n-1}{2}, \\ (H_A^\eta)_{n,j} &= (-1)^{n-j}\sum_{l=0}^{j-1}(-1)^l = \begin{cases} 1 & l \; odd \\ 0 & l \; even \end{cases}.\end{aligned}$$

It is clear that $(H_A^\eta)_{k,n+1} = (H_A^\eta)_{k,n}$, $(H_A^\eta)_{n+1,n+1} = (H_A^\eta)_{n,n}$, $(H_A^\eta)_{n,n+1} = (H_A^\eta)_{n+1,n}$, $(H_A^\eta)_{n+1,j} = (H_A^\eta)_{n,j}$.

So the matrix H_A^η is

$$H_A^\eta = \begin{pmatrix} 2 & 0 & \cdots & 2 & 0 & 1 & 1 \\ 0 & 0 & \cdots & 0 & 0 & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots \\ 2 & 0 & \cdots & 2 & 0 & 1 & 1 \\ 0 & 0 & \cdots & 0 & 0 & 0 & 0 \\ 1 & 0 & \cdots & 1 & 0 & \frac{n+1}{2} & -\frac{n-1}{2} \\ 1 & 0 & \cdots & 1 & 0 & -\frac{n+1}{2} & \frac{n+1}{2} \end{pmatrix},$$

and the kernel is given by

$$\langle e_{2k-1} - e_1,\, e_{2k}, (e_n + e_{n+1}) - e_1 | k \leq \frac{n-1}{2} \rangle.$$

12.2.3. *$Q = D_{n+1}$, n even.* Since $F = \{1, \dots, n-1\}$, we work only with $e_k A e_j$ for $j, k \leq n-1$, and we have to work with a modified basis, so that they are all eigenvectors of η:

For $\underline{k \leq j \leq n-1}$,

$$\begin{aligned} B_{k,j} \;=\; & \{(a_{k-1}a_{k-1}^*)^l a_k^* \cdots a_{j-1}^* | 0 \leq l \leq \min\{k-1, n-j-1\}\} \cup \\ & \{(a_{k-1}a_{k-1}^*)^l a_k^* \cdots (a_{n-1}^* a_{n-1} - a_n^* a_n) a_{n-2} a_j | 0 \leq l \leq k-1\} \cup \\ & \{(a_{k-1}a_{k-1}^*)^l a_k^* \cdots (a_{n-1}^* a_{n-1} + a_n^* a_n) a_{n-2} a_j | 0 \leq l \leq k-1+j-n\}. \end{aligned}$$

For $\underline{j < k \leq n-1}$,

$$\begin{aligned} B_{k,j} \;=\; & \{a_{k-1} \cdots a_j (a_j^* a_j)^l | 0 \leq l \leq \min\{n-k-1, j-1\}\} \cup \\ & \{a_k^* \cdots a_{n-2}^* (a_{n-1}^* a_{n-1} - a_n^* a_n) a_{n-2} \cdots a_j (a_j^* a_j)^l | 0 \leq l \leq j-1\} \cup \\ & \{a_k^* \cdots a_{n-2}^* (a_{n-1}^* a_{n-1} + a_n^* a_n) a_{n-2} \cdots a_j (a_j^* a_j)^l | 0 \leq l \leq j-1+k-n\}. \end{aligned}$$

For $k \leq j \leq n-1$:

$$\begin{aligned} (H_A^\eta)_{k,j} &= \begin{cases} \sum\limits_{l=0}^{k-1} (-1)^l + (-1)^{n-j-1} \sum\limits_{l=0}^{k-1} (-1)^l & k \leq n-j \\ \sum\limits_{l=0}^{n-j-1} (-1)^l + (-1)^{n-j-1} \sum\limits_{l=0}^{k-1} (-1)^l & \\ \quad +(-1)^{n-j} \sum\limits_{l=0}^{k-1+j-n} (-1)^l & k > n-j \end{cases} \\ &= \begin{cases} 2 & k, l \, \mathit{odd} \\ 0 & \mathit{else} \end{cases}. \end{aligned}$$

For $j < k \leq n-1$:

$$\begin{aligned}(H_A^{\eta})_{k,j} &= \left\{\begin{array}{ll} (-1)^{k-j}\sum\limits_{l=0}^{j-1}(-1)^l + (-1)^{n-j-1}\sum\limits_{l=0}^{j-1}(-1)^l & j \leq n-k \\ (-1)^{k-j}\sum\limits_{l=0}^{n-k-1}(-1)^l + (-1)^{n-j-1}\sum\limits_{l=0}^{j-1}(-1)^l & \\ \quad +(-1)^{n-j}\sum\limits_{l=0}^{k-1+j-n}(-1)^l & j > n-k \end{array}\right. \\ &= \left\{\begin{array}{ll} 2 & k, l\, odd \\ 0 & else \end{array}\right. .\end{aligned}$$

So the matrix H_A^{η} is

$$H_A^{\eta} = \begin{pmatrix} 2 & 0 & \cdots & 2 & 0 & 2 \\ 0 & 0 & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 2 & 0 & \cdots & 2 & 0 & 2 \\ 0 & 0 & \cdots & 0 & 0 & 0 \\ 2 & 0 & \cdots & 2 & 0 & 2 \end{pmatrix},$$

and we get immediately its kernel

$$\langle e_{2k+1} - e_1, e_{2k} | 1 \leq k \leq \frac{n}{2} \rangle.$$

12.2.4. $Q = E_7$. We don't use an explicit basis of A here. All we have to know is the number of no-star letters in the monomial basis elements which can be directly obtained from the Hilbert series $H_A(t)$ in the following way: given a monomial x of length l in $e_k A e_j$, n_{kj} the number of arrows in Q on the shortest path from j to k of length $d(k,j)$, x contains $n_{k,j} + \frac{l-d(k,j)}{2}$ arrows in Q.

So we obtain the formula

$$(H_A^{\eta})_{k,j} = (-1)^{n}_{k,j} \left.\frac{H_A(t)_{k,j}}{t^{d(k,j)}}\right|_{t=\sqrt{(-1)}}$$

and compute

$$H_A^{\eta} = \begin{pmatrix} 3 & 0 & 3 & 0 & 0 & 0 & -3 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 3 & 0 & 3 & 0 & 0 & 0 & -3 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -3 & 0 & -3 & 0 & 0 & 0 & 3 \end{pmatrix},$$

and its kernel is

$$\langle e_1 + e_7, e_2, e_3 + e_7, e_4, e_5, e_6 \rangle.$$

12.3. **Result.** Now we give explicit bases for each quiver where $\psi_i \in U^*[-2]$ satisfy the properties given in section 12.1 and $\varepsilon_i \in Y^*[-h-2]$ are taken from 12.2.

Note the duality $HH^6(A) = (HH^5(A))^*$ which was established in [EE2], $\phi_0(z_0) \in U[-2h-2]$, $\varphi_0(\omega_i) \in Y[-h-2]$. We choose ψ_0 such that $\psi_0(\varphi_0(z_0)) = 1$ (from that follows $\psi_k(\varphi_0(z_k)) = z_k\psi_k(\varphi_0(z_0)) = \psi_0(\varphi_0(z_0)) = 1$ and ε_i such that $\varepsilon_i(\phi_0(\omega_j)) = \delta_{ij}$.

12.3.1. $Q = D_{n+1}$, *n odd.* We define

$$\psi_{4k} = [(a^*_{n-1}a_{n-1}a^*_n a_n)^{\frac{n-1}{2}-k}],$$

$$\varepsilon_{2k-1} = [e_{2k-1} - e_1],\ \varepsilon_{2k} = [e_{2k}], \varepsilon_n = [(e_n + e_{n+1}) - e_1], k \leq \frac{n-1}{2}.$$

12.3.2. $Q = D_{n+1}$, *n even.* We define

$$\psi_{4k} = [a^*_{n-1}a_{n-1}(a^*_n a_n a^*_{n-1}a_{n-1})^{\frac{n-2}{2}-k}],$$

$$\varepsilon_{2k+1} = [e_{2k+1} - e_1], \varepsilon_{2k} = [e_{2k}], 1 \leq k \leq \frac{n}{2} - 1.$$

12.3.3. $Q = E_6$. We define

$$\begin{aligned} \psi_0 &= [a^*_3a_3(a^*_2a_2a^*_3a_3)^2], \\ \psi_6 &= [-a^*_3a_3a^*_2a_2], \\ \psi_8 &= [a^*_3a_3 - a^*_2a_2], \end{aligned}$$

$$\varepsilon_3 = [e_3],\ \varepsilon_6 = [e_6].$$

12.3.4. $Q = E_7$. We define

$$\begin{aligned} \psi_0 &= [(a^*_4a_4a^*_3a_3)^4], \\ \psi_8 &= [(a^*_4a_4a^*_3a_3)^2], \\ \psi_{12} &= [a^*_4a_4a^*_3a_3], \end{aligned}$$

$$\varepsilon_1 = [e_1 + e_7], \varepsilon_2 = [e_2], \varepsilon_3 = [e_3 + e_7], \varepsilon_4 = [e_4], \varepsilon_5 = [e_5], \varepsilon_6 = [e_6].$$

12.3.5. $Q = E_8$. We define

$$\begin{aligned} \psi_0 &= [(a^*_4a_4a^*_3a_3)^7], \\ \psi_{12} &= [(a^*_4a_4a^*_3a_3)^4], \\ \psi_{20} &= [(a^*_4a_4a^*_3a_3)^2], \\ \psi_{24} &= [a^*_4a_4a^*_3a_3], \end{aligned}$$

$$\varepsilon_1 = [e_1], \varepsilon_2 = [e_2], \varepsilon_3 = [e_3], \varepsilon_4 = [e_4], \varepsilon_5 = [e_5], \varepsilon_6 = [e_6], \varepsilon_7 = [e_7], \varepsilon_8 = [e_8].$$

13. $HH^6(A)$

$HH^6(A) = U[-2h-2] \oplus Y[-h-2] = HH^0(A)/Im(d_6^*)$, and $Im(d_6^*)$ is spanned by the columns of the matrices H_A^η which were computed in the previous section.

This gives us the following result:

Proposition 13.0.1. *$HH^6(A)$ is a quotient of $HH^0(A)$. Im particular,*

$$HH^6(A) = \begin{cases} HH^0(A) & Q = E_6,\, E_8 \\ HH^0(A)/(\sum\limits_{\substack{i=1 \\ odd}}^{n-2} \omega_i = 0, \omega_n = \omega_{n+1}) & Q = D_{n+1},\, n \text{ odd} \\ HH^0(A)/(\sum\limits_{\substack{i=1 \\ odd}}^{n-1} \omega_i = 0), & Q = D_{n+1},\, n \text{ even} \\ HH^0(A)/(\omega_1 + \omega_3 - \omega_7 = 0) & Q = E_7 \end{cases}.$$

14. Products involving $HH^0(A) = Z$

Recall the decomposition $HH_0(A) = \mathbb{C} \oplus (U[-2])_+ \oplus L[h-2]$. It is clear that the $\mathbb{C}$-part acts on $HH^i(A)$ as the usual multiplication with $\mathbb{C}$, with z_0 as identity. From the periodicity of the Schofield resolution with period 6, it follows that the multiplication with $\varphi(z_0) \in HH^6(A)$ gives the natural isomorphism $HH^i(A) \to HH^{i+6}(A)$ for $i \geq 1$.

We summarize all products not involving the $\mathbb{C}$-part.

14.1. $HH^0(A) \times HH^0(A) \twoheadrightarrow HH^0(A)$.

This is already done in the $HH^0(A)$-section of this paper. We state the results:

14.1.1. $Q = D_{n+1}$, *n odd.* The products are

$$z_{4j} z_{4k} = \begin{cases} z_{4(j+k)} & j+k < \frac{n-1}{2} \\ \omega_n - \omega_{n+1} & j+k = \frac{n-1}{2} \\ 0 & j+k > \frac{n-1}{2} \end{cases}.$$

14.1.2. $Q = D_{n+1}$, *n even.* The products are

$$z_{4j} z_{4k} = \begin{cases} z_{4(j+k)} & j+k < \frac{n-1}{2} \\ 0 & j+k \geq \frac{n-1}{2} \end{cases}.$$

14.1.3. E_6. All products are zero.

14.1.4. E_7. The only nonzero product is $z_8^2 = \omega_1 + \omega_3 - \omega_7$.

14.1.5. E_8. The only nonzero product is $z_{12}^2 = z_{24}$.

14.2. $HH^0(A) \times HH^1(A) \twoheadrightarrow HH^1(A)$.

From the definition of the maps θ_k (which are generated by the central elements z_k), it follows that the Z-action is natural, i.e. the multiplication rule is the same as with the z_k counterpart: $z_k \theta_0 = \theta_k$.

We state the other nonzero products:

14.2.1. $Q = D_{n+1}$. We have $z_{4j}\theta_{4k} = \theta_{4(j+k)}$ if $j + k < \frac{n-1}{2}$.

14.2.2. E_8. We have $z_{12}\theta_{12} = \theta_{24}$.

14.3. $HH^0(A) \times HH^i(A) \twoheadrightarrow HH^i(A)$**,** $i = 2$ **or** 3**.** $HH^2(A) = K[-2]$ and $HH^3(A) = K^*[-2]$ live in only one degree, so $(U[-2])_+$ $HH^0(A)$ acts by zero.

14.4. $HH^0(A) \times HH^4(A) \twoheadrightarrow HH^4(A)$**.** We defined ζ_k, such that $z_k\zeta_k = \zeta_0$ holds. By degree arguments, only these other products are nonzero:

14.4.1. $Q = D_{n+1}$. For $l < k$, $z_{4l}\zeta_{4k} = \zeta_{4(k-l)}$ (since $z_{4(k-l)}(z_{4l}\zeta_{4k}) = (z_{4(k-l)}z_{4l})\zeta_{4k}) = \zeta_0$, and $\zeta_{4(k-l)}$ is only one element of degree $-4 - 4(k - l)$ in $HH^4(A)$).

14.4.2. $Q = E_8$. We have $z_{12}\zeta_{24} = \zeta_{12}$ (since $z_{12}(z_{12}\zeta_{24}) = (z_{12}z_{12})\zeta_{24} = \zeta_0$, and ζ_{12} is the only element of degree -16 in $HH^4(A)$).

14.5. $HH^0(A) \times HH^5(A) \twoheadrightarrow HH^5(A)$**.** By definition, $z_k\psi_k = \psi_0$ holds. Since $\psi_i \in U^*[-2]$ corresponds to $\zeta_i \in U^*[-2]$ in $HH^4(A)$ with the rule $z_k\psi_k = \psi_0$ corresponding to $z_k\zeta_k = \zeta_0$ above, the multiplication rules of ψ_k with elements in $HH^0(A)$ can be derived from above.

Products involving $\omega_i \in L[h-2] \subset HH^0(A)$ and $\varepsilon_j = \sum_{k\in F} \lambda_k e_k \in Y^*[-h-2]$ are easy to calculate: $\omega_i\varepsilon_j = \lambda_i[\omega_i] = \lambda_i\psi_0$.

Proposition 14.5.1. *The multiplication* $((U[-2])_+)\times Y^*[-h-2] \to HH^5(A)$ *is zero*

We will show this for any quiver separately.

14.5.1. $Q = D_{n+1}$, *n odd.* For $l < k$, $z_{4l}\psi_{4k} = \psi_{4(k-l)}$.

The nonzero products involving $\omega_i \in L[h-2] \subset HH^0(A)$ and $\varepsilon_j \in Y^*[-h-2]$ are

$$\omega_{2k-1}\varepsilon_{2k-1} = \omega_{2k}\varepsilon_{2k} = \omega_n\varepsilon_n = \omega_{n+1}\varepsilon_n = \omega_1\varepsilon_{2k-1} = \omega_1\varepsilon_n = \psi_0,$$

$$\omega_1\varepsilon_{2k-1} = \omega_1\varepsilon_n = -\psi_0.$$

We show $(U[-2])_+ \times Y^*[-h-2] \xrightarrow{0} HH^5(A)$: by degree argument, $z_{4k}\varepsilon_i = \lambda\psi_{2n-2-4k}$. Then $z_{2n-2-4k}(z_{4k}\varepsilon_i) = \lambda z_{2n-2-4k}\psi_{2n-2-4k} = \lambda\psi_0$, and by associativity this equals $(z_{2n-2-4k}z_{4k})\varepsilon_i = (\omega_n - \omega_{n+1})\varepsilon_i = 0$, so $\lambda = 0$.

14.5.2. $Q = D_{n+1}$, *n even.* For $l < k$, $z_{4l}\psi_{4k} = \psi_{4(k-l)}$.

The nonzero products involving $\omega_i \in L[h-2] \subset HH^0(A)$ and $\varepsilon_j \in Y^*[-h-2]$ are

$$\omega_{2k+1}\varepsilon_{2k+1} = \omega_{2k}\varepsilon_{2k} = \psi_0,$$
$$\omega_1\varepsilon_{2k+1} = -\psi_0.$$

We show $(U[-2])_+ \times Y^*[-h-2] \xrightarrow{0} HH^5(A)$: by degree argument, $z_{4k}\varepsilon_i = \lambda\psi_{2n-2-4k}$. Then $z_{2n-2-4k}(z_{4k}\varepsilon_i) = \lambda z_{2n-2-4k}\psi_{2n-2-4k} = \lambda\psi_0$, and this equals $(z_{2n-2-4k}z_{4k})\varepsilon_i = 0$, so $\lambda = 0$.

14.5.3. $Q = E_6$. The nonzero products involving $\omega_i \in L[h-2] \subset HH^0(A)$ and $\varepsilon_j \in Y^*[-h-2]$ are

$$\omega_3\varepsilon_3 = \omega_6\varepsilon_6 = \psi_0.$$

By degree argument, $(U[-2])_+ \times Y^*[-h-2] \xrightarrow{0} HH^5(A)$.

14.5.4. $Q = E_7$. The nonzero products involving $\omega_i \in L[h-2] \subset HH^0(A)$ and $\varepsilon_j \in Y^*[-h-2]$ are

$$\omega_1\varepsilon_1 = \omega_2\varepsilon_2 = \omega_3\varepsilon_3 = \omega_4\varepsilon_4 = \omega_5\varepsilon_5 = \omega_6\varepsilon_6 = \omega_7\varepsilon_1 = \omega_7\varepsilon_3 = \psi_0.$$

We show $(U[-2])_+ \times Y^*[-h-2] \xrightarrow{0} HH^5(A)$: by degree argument, only products involving z_8 may eventually be nontrivial,

$$z_8\varepsilon_i = \lambda\psi_8, \quad \lambda \in \mathbb{C}.$$

Then

$$z_8(z_8\varepsilon_i) = \lambda z_8\psi_8 = \lambda\psi_0,$$

and by associativity this equals

$$z_8^2\varepsilon_i = (\omega_1 + \omega_3 - \omega_7)\varepsilon_i = 0,$$

so $\lambda = 0$.

15. Products involving $HH^1(A)$

15.1. $HH^1(A) \times HH^1(A) \xrightarrow{0} HH^2(A)$**.** This follows by degree argument since $\deg HH^1(A) > 0$, $\deg HH^2(A) = -2$.

15.2. $HH^1(A) \times HH^2(A) \twoheadrightarrow HH^3(A)$**.** $HH^2(A)$ and $HH^3(A)$ are trivial for $Q = D_{n+1}$ where n is odd and for $Q = E_7$, E_8.

We know that $HH^1(A)$ is generated by maps θ_k and $HH^2(A)$ by f_i ($i \neq \nu(i)$, and we lift

$$\begin{aligned} f_i : A \otimes A[2] &\longrightarrow A, \\ 1 \otimes 1 &\longmapsto e_i - e_{\nu(i)} \end{aligned}$$

to

$$\begin{aligned} \hat{f}_i : A \otimes A[2] &\longrightarrow A \otimes A, \\ 1 \otimes 1 &\longmapsto e_i \otimes e_i - e_{\nu(i)} \otimes e_{\nu(i)}. \end{aligned}$$

Then

$$\hat{f}_i d_3(1 \otimes 1) = \hat{f}_i(\sum_{x_j \in B} x_j \otimes x_j^*) = \sum_{x_j \in B} x_j e_i \otimes e_i x_j^* - x_j e_{\nu(i)} \otimes e_{\nu(i)} x_j^*.$$

To compute the lift Ωf_i, we need to find out the preimage of $\sum x_j e_i \otimes e_i x_j^* - x_j e_{\nu(i)} \otimes e_{\nu(i)} x_j^*$ under d_1.

Definition 15.2.1. *Let $b_1, \dots, b_k$ be arrows, p the monomial $\pm b_1 \cdots b_k$ and define*

$$v_p := \pm(1 \otimes b_1 \otimes b_2 \cdots b_k + b_1 \otimes b_2 \otimes b_3 \cdots b_k + \ldots + b_1 \cdots b_{k-1} \otimes b_k \otimes 1),$$

and for $i < j$,

$$v_p^{(i,j)} := \pm \sum_{l=i}^{j} b_1 \cdots b_{l-1} \otimes b_l \otimes b_{l+1} \cdots b_k$$

We will use the following lemma in our computations.

Lemma 15.2.2. *In the above setting,*

$$d_1(v_p) = \pm(b_1 \cdots b_k \otimes 1 - 1 \otimes b_1 \cdots b_k).$$

From that, we see immediately that when assuming all x_j are monomials (what we can do), then

$$\hat{f}_i(\sum_{x_j \in B} x_j \otimes x_j^*) = d_1(\sum_{x_j \in B} v_{x_j e_i x_j^*}^{(1,\deg(x_j))} - v_{x_j e_{\nu(i)} x_j^*}^{(1,\deg(x_j))}) + 1 \otimes \underbrace{\sum_{x_j \in B} (x_j e_i x_j^* - x_j e_{\nu(i)} x_j^*)}_{=0},$$

so we have

$$\begin{aligned} \Omega f_i : \Omega^3(A) &\longrightarrow \Omega(A), \\ 1 \otimes 1 &\longmapsto \sum_{x_j \in B} v_{x_j e_i x_j^*}^{(1,\deg(x_j))} - v_{x_j e_{\nu(i)} x_j^*}^{(1,\deg(x_j))}. \end{aligned}$$

Then

$$\theta_k(\sum_{x_j \in B} v_{x_j e_i x_j^*}^{(1,\deg(x_j))} - v_{x_j e_{\nu(i)} x_j^*}^{(1,\deg(x_j))}) = z_k(\sum_{x_j \in B_{-,i}} s(x_j) x_j x_j^* - \sum_{x_j \in B_{-,\nu(i)}} s(x_j) x_j x_j^*),$$

where for $s(x_j)$ is the number of arrows in Q^* in the monomial expression of x_j.

So we get

$$(\theta_k \circ \Omega f_i)(1 \otimes 1) = z_k(\sum_{x_j \in B_{-,i}} s(x_j) x_j x_j^* - \sum_{x_j \in B_{-,\nu(i)}} s(x_j) x_j x_j^*).$$

Under our identification in [EE2, Subsection 4.5.],

$$\theta_k f_i = [z_k(\sum_{l \in I} \sum_{x_j \in B_{l,i}} s(x_j) \omega_l - \sum_{l \in I} \sum_{x_j \in B_{l,\nu(i)}} s(x_j) \omega_l)] \in HH^3(A).$$

All products are zero if z_k lies in a positive degree, so we only have to calculate the products where $k = 0$.

We make the following

Proposition 15.2.3. *The multiplication with θ_0 induces a symmetric isomorphism*

$$\alpha : HH^2(A) = K[-2] \overset{\cong}{\to} K^*[-2] = HH^3(A).$$

Now we have to work with explicit basis elements $x_j \in Ae_i$, $i \neq \nu(i)$, so we treat the Dynkin quivers separately and find the matrix M_α which represents this map.

15.2.1. $Q = D_{n+1}$, *n even.* We can work with the basis given in section 6 and compute

$$\theta_0 f_n = \frac{n}{2}([\omega_{n+1}] - [\omega_n]) = -nh_n \tag{15.2.4}$$

because of the relation $[\omega_n] + [\omega_{n+1}] = 0$ in $HH^3(A)$. α is given by the matrix

$$M_\alpha = (-n).$$

15.2.2. E_6. We will write out the basis elements of Ae_1, Ae_5:

$$B_{1,1} = \langle e_1, a_1a_2a_5^*a_5a_2^*a_1^*\rangle,$$
$$B_{2,1} = \langle a_1^*, a_2a_5^*a_5a_2^*a_1^*, a_2a_3^*a_3a_5^*a_5a_2^*a_1^*\rangle,$$
$$B_{3,1} = \langle a_2^*a_1^*, a_3^*a_3a_2^*a_1^*, a_3^*a_3a_3^*a_3a_2^*a_1^*, a_5^*a_5a_3^*a_3a_3^*a_3a_2^*a_1^*\rangle,$$
$$B_{4,1} = \langle a_3a_2^*a_1^*, a_3a_5^*a_5a_2^*a_1^*, a_3a_5^*a_5a_3^*a_3a_5^*a_5a_2^*a_1^*\rangle,$$
$$B_{5,1} = \langle a_4a_3a_2^*a_1^*, a_4a_3a_5^*a_5a_3^*a_3a_5^*a_5a_2^*a_1^*\rangle,$$
$$B_{6,1} = \langle a_5a_2^*a_1^*, a_5a_3^*a_3a_5^*a_5a_2^*a_1^*\rangle,$$

and

$$e_iAe_5 = \langle \eta(x) | x \in e_{\nu(i)}Ae_1\rangle,$$

where $\eta(a) = -\epsilon_a\bar{a}$ and for any arrow $a : i \to j$, $\bar{a}$ is the arrow $j :\to i$, so η preserves the number of star letters of a monomial x. From this, we obtain

$$\theta_0 f_1 = -4[\omega_1] - 2[\omega_2] + 2[\omega_4] + 4[\omega_5] = -8h_1 - 4h_2.$$

because of the relations $\omega_1 + \omega_4 = \omega_2 + \omega_3 = 0$ in $HH^3(A)$.

We do the same thing for Ae_2 and Ae_4:

$$B_{1,2} = \langle a_1, a_1a_2a_5^*a_5a_2^*, a_1a_2a_5^*a_5a_3^*a_3a_2^*\rangle,$$
$$B_{2,2} = \langle e_2, a_2a_2^*, a_2a_5^*a_5a_2^*, a_2a_3^*a_3a_5^*a_5a_2^*, a_2a_5^*a_5a_3^*a_3a_2^*, a_2a_5^*a_5a_3^*a_3a_5^*a_5a_2^*\rangle,$$
$$\begin{aligned} B_{3,2} = \langle & a_2^*, a_5^*a_5a_2^*, a_3^*a_3a_2^*, a_5^*a_5a_3^*a_3a_2^*, a_3^*a_3a_5^*a_5a_2^*, \\ & a_3^*a_3a_5^*a_5a_3^*a_3a_2^*, a_5^*a_5a_3^*a_3a_5^*a_5a_2^*, a_5^*a_5a_3^*a_3a_5^*a_5a_3^*a_3a_2^*\rangle, \end{aligned}$$
$$\begin{aligned} B_{4,2} = \langle & a_3a_2^*, a_3a_5^*a_5a_2^*, a_3a_3^*a_3a_2^*, a_3a_3^*a_3a_5^*a_5a_2^*, \\ & a_3a_5^*a_5a_3^*a_3a_5^*a_5a_2^*, a_3a_3^*a_3a_5^*a_5a_3^*a_3a_5^*a_5a_2^*\rangle, \end{aligned}$$
$$B_{5,2} = \langle a_4a_3a_2^*, a_4a_3a_5^*a_5a_2^*, a_4a_3a_5^*a_5a_3^*a_3a_5^*a_5a_2^*\rangle,$$
$$B_{6,2} = \langle a_5a_2^*, a_5a_3^*a_3a_2^*, a_5a_3^*a_3a_5^*a_5a_2^*, a_5a_3^*a_3a_5^*a_5a_3^*a_3a_2^*\rangle,$$

and we get the basis elements for e_iAe_3 from $\eta(x_j)$ where $x_j \in e_{\nu(i)}Ae_4$. Since η preserves the number of star-letters of a monomial, we can immediately calculate

$$\theta_0 f_2 = -2[\omega_1] - 4[\omega_2] + 4[\omega_4] + 2[\omega_5] = -4h_1 - 8h_2$$

because of the relations $[\omega_1] + [\omega_4] = [\omega_2] + [\omega_3] = 0$ in $HH^3(A)$.

So α is given by the symmetric, nondegenerate matrix

$$M_\alpha = \begin{pmatrix} -8 & -4 \\ -4 & -8 \end{pmatrix}. \tag{15.2.5}$$

15.3. $HH^1(A) \times HH^3(A) \xrightarrow{0} HH^4(A)$**.** This follows by degree argument: $\deg HH^1(A) \geq 0$, $\deg HH^3(A) = -2$, but $\deg HH^4(A) \leq -4$.

15.4. $HH^1(A) \times HH^4(A) \to HH^5(A)$**.**

Proposition 15.4.1. *Given $\theta_k \in HH^1(A)$ and $\zeta_l \in HH^4(A)$, we get the following cup product:*

$$\theta_k \zeta_l = \psi_l z_k. \tag{15.4.2}$$

Proof. It is enough to show $\theta_0\zeta_0 = \psi_0$: $z_l(\theta_0\zeta_l) = \theta_0\zeta_0\psi_0$ implies that $(\theta_0\zeta_l) = \psi_l$, and the equation above follows from $\theta_k = z_k\theta_0$.

Let in general $x = \sum_{a \in \bar{Q}} a \otimes x_a \in HH^4(A)$. Then x represents the map

$$\begin{aligned} x := A \otimes V \otimes \mathcal{N}h &\longrightarrow A, \\ 1 \otimes a_i \otimes 1 &\longmapsto -x_{a_i^*} \\ 1 \otimes a_i^* \otimes 1 &\longmapsto x_{a_i}, \end{aligned}$$

and it lifts to

$$\begin{aligned} \hat{x} : A \otimes V \otimes \mathcal{N}[h] &\longrightarrow A \otimes A, \\ 1 \otimes a \otimes 1 &\longmapsto -1 \otimes x_{a^*} \\ 1 \otimes a^* \otimes 1 &\longmapsto 1 \otimes x_a. \end{aligned}$$

Then

$$\begin{aligned} (\hat{x} \circ d_5)(1 \otimes 1) &= \hat{x}(\sum_{a \in \bar{Q}} \epsilon_a a \otimes a^* \otimes 1 + \sum_{a \in \bar{Q}} \epsilon_a 1 \otimes a \otimes a^*) \\ &= \sum_{a \in Q} a \otimes x_a - \sum_{a \in Q} 1 \otimes x_a \eta(a) + \sum_{a \in Q} a^* \otimes x_{a^*} - \sum_{a \in Q} 1 \otimes x_{a^*}\eta(a^*) \\ &= \sum_{a \in Q} a \otimes x_a - \sum_{a \in Q} 1 \otimes a x_a \sum_{a \in Q} a^* \otimes x_{a^*} - \sum_{a \in Q} 1 \otimes a^* x_{a^*} \\ &= d_1(\sum_{a \in Q} 1 \otimes a \otimes x_a + 1 \otimes a^* \otimes x_{a^*}), \end{aligned}$$

so we have

$$\begin{array}{rcl} \Omega x : \Omega^5(A) & \longrightarrow & \Omega(A), \\ 1 \otimes 1 & \longmapsto & \sum\limits_{a \in Q} 1 \otimes a \otimes x_a + 1 \otimes a^* \otimes x_{a^*}, \end{array}$$

and this gives us

$$(\theta_0 \circ x)(1 \otimes 1) = \sum_{a \in Q} a^* x_{a^*},$$

so the cup product is

$$\theta_0 \cdot x = \sum_{a \in Q} a^* x_{a^*}. \tag{15.4.3}$$

It can be easily checked by using explicit elements that the RHS is ψ_0 for $x = \zeta_0$, but we the reason here why this is true: for $x = \sum a \otimes x_a = \zeta_0$, the RHS becomes

$$\sum_{a \in Q} a^* x_{a^*} = \sum_{a \in Q} (a^*, x_{a^*})[\omega_{t(a)}],$$

where $(-,-) : A \times A \to \mathbb{C}$ is the bilinear form attached to A as a Frobenius algebra (see 5.3).

But under the bilinear form on $V \otimes A$, given in [EE2, Subsection 4.3.] which induces the duality $HH^4(A) = (HH^1(A))^*$,

$$(a \otimes x_a, b \otimes x_b) = \delta_{a,b^*} \epsilon_a (x_a, x_b),$$

$$\sum_{a \in Q} (a^*, x_{a^*}) = (\theta_0, \zeta_0) = 1.$$

So for $x = \zeta_0$, equation (15.4.3) becomes

$$\theta_0 \zeta_0 = (\theta_0, \zeta_0)\psi_0 = \psi_0, \tag{15.4.4}$$

because $[\omega_i] = \psi_0$ in $HH^5(A)$ for all $i \in I$. □

15.5. $HH^1(A) \times HH^5(A) \to HH^6(A)$**.** We know that

$$\begin{array}{rcl} 0 & \leq & \deg(HH^1(A)) \leq h - 4, \\ -h - 2 & \leq & \deg(HH^5(A)) \leq -2, \\ -2h & \leq & \deg(HH^6(A)) \leq -h - 2, \end{array}$$

so the product is trivial unless we pair the lowest degree parts of $HH^1(A)$ (generated by θ_0) and $HH^5(A)$ (which is $Y^*[-h-2]$). The product will then live in degree $-h - 2$ which is the top degree part of $HH^6(A)$, the space $Y[-h - 2]$.

Given an element $\psi \in HH^5(A)(-h - 2))$ which has the form

$$\begin{array}{rcl} \psi : A \otimes \mathcal{N}[h + 2] & \longrightarrow & A, \\ 1 \otimes 1 & \longmapsto & \sum\limits_{i \in F} \lambda_i e_i \in R, \end{array}$$

this lifts to

$$\begin{array}{rcl}\hat{\psi}: A\otimes\mathcal{N}[h+2] & \longrightarrow & A\otimes A,\\ 1\otimes 1 & \longmapsto & \sum\limits_{i\in F}\lambda_i e_i\otimes e_i.\end{array}$$

Then

$$\begin{aligned}\hat{\psi}(d_6(1\otimes 1)) &= \hat{\psi}(\sum_{x_j\in B}\sum_{i\in F} x_j\otimes x_j^*) = \hat{\psi}(\sum_{x_j\in B}\sum_{i\in F}\eta(x_j)\otimes\eta(x_j^*))\\ &= \sum_{x_j\in B}\sum_{i\in F}\lambda_i\eta(x_j)e_i\otimes e_i x_j^*\\ &= d_1(\sum_{i\in F}\sum_{x_j\in B}\lambda_i v^{(1,\deg(x_j))}_{\eta(x_j)e_i x_j^*}) + 1\otimes\underbrace{\sum_{i\in F}\lambda_i\eta(x_j)e_i x_j^*}_{=0},\end{aligned}$$

so ψ lifts to

$$\begin{array}{rcl}\Omega\psi:\Omega^6(A) & \longrightarrow & \Omega(A),\\ 1\otimes 1 & \longmapsto & \sum\limits_{i\in F}\sum\limits_{x_j\in B}\lambda_i v^{(1,\deg(x_j))}_{\eta(x_j)e_i x_j^*}.\end{array}$$

We get

$$(\theta_0\circ\Omega\psi)(1\otimes 1) = \sum_{i\in F}\sum_{x_j\in Ae_i}\lambda_i s(x_j)\eta(x_j)x_j^*,$$

where $s(x_j)$ is the number of arrows in Q^* in the monomial expression of x_j (or in general if x_j is a homogeneous polynomial where each monomial term has the same number of arrows in Q^*, then $s(x_j)$ is the number of Q^*-arrows in each monomial term).

Under our identifications in [EE2, Subsection 4.5.],

$$\theta_0\psi = \sum_{i\in F}\sum_{x_j\in Ae_i}\lambda_i s(x_j)\eta(x_j)x_j^* = \sum_{i,k\in F}\sum_{x_j\in e_kAe_i}\lambda_i s(x_j)\eta(x_j)x_j^*$$

To simplify this computation, we will choose a basis, such that all $x_j\in e_kAe_l$ for some $k,l\in I$ and that additionally x_j is an eigenvector of η for $k,l\in F$ (since η is an involution on e_kAe_l for $k,l\in F$). Let $B^+_{k,l}$ be a basis of $(e_kAe_l)_+=\ker(\eta|_{e_kAe_l}-1)$ and $B^-_{k,l}$ a basis of $(e_kAe_l)_-=\ker(\eta|_{e_kAe_l}+1)$.

Let us define

$$\kappa_{k,l} = \sum_{x_j\in B^+_{k,l}} s(x_j) - \sum_{x_j\in B^-_{k,l}} s(x_j). \tag{15.5.1}$$

Then the above equation becomes

$$\theta_0\psi = \sum_{l\in F}\lambda_l\sum_{k\in F}\kappa_{k,l}\varphi_0(\omega_k). \tag{15.5.2}$$

Proposition 15.5.3. *The multiplication by θ_0 induces a skew-symmetric isomorphism*

$$\beta : Y^*[-h-2] \overset{\cong}{\to} Y[-h-2].$$

We will treat the Dynkin quivers separately and find the matrix M_β which represents β for each of these quivers.

15.5.1. $Q = D_{n+1}$, *n odd.* We use the same basis as given in section 6. Recall that these basis elements have the property $\eta(x) = (-1)^{n_x}x$ where n_x is the number of Q-arrows in the monomial expression of x.

We can compute that for $k \le l \le n-1$,

$$\begin{aligned}\kappa_{k,l} &= \begin{cases} \sum_{j=0}^{k-1}(-1)^j(l-k+j)+(-1)^{n-l}\sum_{j=0}^{k-1}(-1)^j(n-k+j), & k \le n-l \\ \sum_{j=0}^{n-l-1}(-1)^j(l-k+j)+(-1)^{n-l}\sum_{j=0}^{k-1}(-1)^j(n-k+j) & \\ +(-1)^{n-l}\sum_{j=0}^{k-1+l-n}(-1)^j(n-k+j), & k > n-l \end{cases} \\ &= \begin{cases} n-k+l-1 & k\,\mathit{odd}, \quad l\,\mathit{odd} \\ l-n & k\,\mathit{odd}, \quad l\,\mathit{even} \\ -k & k\,\mathit{even}, \quad l\,\mathit{odd} \\ 0 & k\,\mathit{even}, \quad l\,\mathit{even} \end{cases}\end{aligned}$$

for $l \le k \le n-1$,

$$\begin{aligned}\kappa_{k,l} &= \begin{cases} (-1)^{k-l}\sum_{j=0}^{l-1}(-1)^j j+(-1)^{n-l}\sum_{j=0}^{l-1}(-1)^j(n-k+j) & j \le n-k \\ (-1)^{k-l}\sum_{j=0}^{n-k-1}(-1)^j j+(-1)^{n-l}\sum_{j=0}^{l-1}(-1)^j(n-k+j) & \\ +(-1)^{n-l}\sum_{j=0}^{l+k-n-1}(-1)^j(n-k+j) & j > n-k \end{cases} \\ &= \begin{cases} n-k+l-1 & k\,\mathit{odd}, \quad l\,\mathit{odd} \\ l & k\,\mathit{odd}, \quad l\,\mathit{even} \\ n-k & k\,\mathit{even}, \quad l\,\mathit{odd} \\ 0 & k\,\mathit{even}, \quad l\,\mathit{even} \end{cases}\end{aligned}$$

for $k, l \leq n-1$,

$$\begin{aligned}
\kappa_{k,n} = \kappa_{k,n+1} &= \sum_{j=0}^{k-1}(-1)^j(n-k-j) = \begin{cases} n - \frac{k+1}{2} & k\, odd \\ -\frac{k}{2} & k\, even \end{cases} \\
\kappa_{n,l} = \kappa_{n+1,l} &= (-1)^{n-j}\sum_{j=0}^{l-1}(-1)^j j = \begin{cases} n - \frac{l-1}{2} & l\, odd \\ \frac{l}{2} & k\, even \end{cases} \\
\kappa_{n,n} = \kappa_{n+1,n+1} &= \sum_{j=0}^{\frac{n-1}{2}} 2j = \frac{n^2-1}{4} \\
\kappa_{n+1,n} = \kappa_{n,n+1} &= \sum_{j=0}^{\frac{n-3}{2}}(-1\cdot(1+2l)) = -\left(\frac{n-1}{2}\right)^2
\end{aligned}$$

$Y^*[-h-2]$ has basis $\varepsilon_{2k+1} = [e_{2k+1} - e_1]$ $(0 \leq k \leq \frac{n-3}{2})$, $\varepsilon_{2k} = [e_{2k}]$ $(k \leq \frac{n-1}{2})$, $\varepsilon_n = [e_n + e_{n+1} - e_1]$, and we can calculate the products

$$\begin{aligned}
\theta_0\varepsilon_{2k+1} &= \sum_{i\in F}(\kappa_{i,2k+1} - \kappa_{i,1})\varphi_0(\omega_i) \\
&= 2k\sum_{\substack{i=1\\odd}}^{n-2}\varphi_0(\omega_i) - n\sum_{\substack{i=2\\even}}^{2k}\varphi_0(\omega_i) + k\varphi_0(\omega_n + \omega_{n+1}), \\
\theta_0\varepsilon_{2k} &= \sum_{i\in F}(\kappa_{i,2k+1})\varphi_0(\omega_i) \\
&= (2k-n)\sum_{\substack{i=1\\odd}}^{2k-1}\varphi_0(\omega_i) + 2k\sum_{\substack{i=2k+1\\odd}}^{n-2}\varphi_0(\omega_i) + k\varphi_0(\omega_n + \omega_{n+1}), \\
\theta_0\varepsilon_n &= \sum_{i\in F}(\kappa_{i,n} + \kappa_{n+1,1} - \kappa_{i,1})\varphi_0(\omega_i) \\
&= (n-1)\sum_{i=1,odd}^{n-2}\varphi_0(\omega_i) - n\sum_{\substack{i=2\\even}}^{n-1}\varphi_0(\omega_i) + \frac{n-1}{2}\varphi_0(\omega_n + \omega_{n+1})
\end{aligned}$$

We use the defining relations in $Y[-h-2]$,

$$\begin{aligned}
\varphi_0(\omega_1) &= -\varphi_0(\sum_{\substack{i=3\\odd}}^{n-2}\varphi_0(\omega_i) - \varphi_0(\omega_n)) \\
\varphi_0(\omega_{n+1}) &= \varphi_0(\omega_n)
\end{aligned}$$

to write the RHS of the above cup product calculations in terms of the basis $(\omega_i)_{2\leq i\leq n}$:

$$\begin{aligned}
\theta_0\varepsilon_{2k+1} &= -n\sum_{\substack{i=2\\ even}}^{2k}\varphi_0(\omega_i),\\
\theta_0\varepsilon_{2k} &= n\sum_{\substack{i=2k+1\\ odd}}^{n-2}\varphi_0(\omega_i)+n\varphi_0(\omega_n),\\
\theta_0\varepsilon_{n} &= -n\sum_{\substack{i=2\\ even}}^{n-1}\varphi_0(\omega_i).
\end{aligned}$$

β is given by the skew-symmetric, nondegenerate matrix

$$M_\beta=\begin{pmatrix}
0 & -n & 0 & -n & \dots & \dots & -n & 0 & -n & 0 & -n\\
n & 0 & 0 & 0 & \dots & \dots & 0 & 0 & 0 & 0 & 0\\
0 & 0 & 0 & -n & \dots & \dots & -n & 0 & -n & 0 & -n\\
n & 0 & n & 0 & \dots & \dots & 0 & 0 & 0 & 0 & 0\\
0 & 0 & 0 & 0 & \dots & \dots & -n & 0 & -n & 0 & -n\\
\vdots & \vdots & \vdots & \vdots & \ddots & & \vdots & \vdots & \vdots & \vdots & \vdots\\
\vdots & \vdots & \vdots & \vdots & & \ddots & \vdots & \vdots & \vdots & \vdots & \vdots\\
n & 0 & n & 0 & \dots & \dots & 0 & 0 & 0 & 0 & 0\\
0 & 0 & 0 & 0 & \dots & \dots & 0 & 0 & -n & 0 & -n\\
n & 0 & n & 0 & \dots & \dots & 0 & n & 0 & 0 & 0\\
0 & 0 & 0 & 0 & \dots & \dots & 0 & 0 & 0 & 0 & -n\\
n & 0 & n & 0 & \dots & \dots & 0 & n & 0 & n & 0
\end{pmatrix}$$

with respect to the chosen basis $\varepsilon_2,\varepsilon_3,\dots\varepsilon_n$ of $Y^*[-h-2]$ and the dual basis $\varphi_0(\omega_2),\varphi_0(\omega_3)\dots\varphi_0(\omega_n)$ of $Y[-h-2]$).

15.5.2. $Q=D_{n+1}$, *n even.* We use the same basis as in section 12.2.3 for our computations.

For $k\leq l\leq n-1$,

$$\begin{aligned}
\kappa_{k,l} &= \begin{cases}
\sum_{j=0}^{k-1}(-1)^j(l-k+j)+(-1)^{n-l+1}\sum_{j=0}^{k-1}(-1)^j(n-k+j), & k\leq n-l\\
\sum_{j=0}^{n-l-1}(-1)^j(l-k+j)+(-1)^{n-l-1}\sum_{j=0}^{k-1}(-1)^j(n-k+j) & \\
\quad +(-1)^{n-l}\sum_{j=0}^{k-1+l-n}(-1)^j(n-k+j), & k>n-l
\end{cases}\\
&= \begin{cases}
n-k+l-1 & k\,odd,\quad l\,odd\\
l-n & k\,odd,\quad l\,even\\
-k & k\,even,\quad l\,odd\\
0 & k\,even,\quad l\,even
\end{cases}
\end{aligned}$$

for $l \leq k \leq n-1$,

$$\kappa_{k,l} = \begin{cases} (-1)^{k-l}\sum_{j=0}^{l-1}(-1)^j j + (-1)^{n-l+1}\sum_{j=0}^{l-1}(-1)^j(n-k+j) & j \leq n-k \\ (-1)^{k-l}\sum_{j=0}^{n-k-1}(-1)^j j + (-1)^{n-l+1}\sum_{j=0}^{l-1}(-1)^j(n-k+j) & \\ +(-1)^{n-l}\sum_{j=0}^{l+k-n-1}(-1)^j(n-k+j) & j > n-k \end{cases}$$

$$= \begin{cases} n-k+l-1 & k\,odd, \quad l\,odd \\ l & k\,odd, \quad l\,even \\ n-k & k\,even, \quad l\,odd \\ 0 & k\,even, \quad l\,even \end{cases}$$

$Y^*[-h-2]$ has basis $\varepsilon_{2k} = [e_{2k}]$, $\varepsilon_{2k+1} = [e_{2k+1} - e_1]$ $(1 \leq k \leq \frac{n-2}{2})$, and we calculate the products

$$\begin{aligned} \theta_0\varepsilon_{2k+1} &= \sum_{i\in F}(\kappa_{i,2k+1} - \kappa_{i,1})\varphi_0(\omega_i) \\ &= 2k\sum_{\substack{i=1,\\ odd}}^{n-1}\varphi_0(\omega_i) - n\sum_{\substack{i=2\\ even}}^{2k}\varphi_0(\omega_i), \\ \theta_0\varepsilon_{2k} &= \sum_{i\in F}(\kappa_{i,2k})\varphi_0(\omega_i) \\ &= (2k-n)\sum_{\substack{i=1\\ odd}}^{2k-1}[\omega_i] + 2k\sum_{i=2k+1}^{n-2}\varphi_0(\omega_i), \end{aligned}$$

and we use the defining relation of $Y[-h-2]$,

$$\varphi_0(\omega_1) = -\sum_{\substack{i=3\\ odd}}^{n-2}\varphi(\omega_i)$$

to write the results of the cup product calculations in terms of the basis $\varphi_0(\omega_2), \varphi_0(\omega_3), \ldots, \varphi_0(\omega_{n-1})$. We get

$$\begin{aligned} \theta_0\varepsilon_{2k+1} &= -n\sum_{\substack{i=2\\ even}}^{2k}\varphi_0(\omega_i) \\ \theta_0\varepsilon_{2k} &= n\sum_{\substack{i=2k+1\\ odd}}^{n-1}\varphi_0(\omega_i). \end{aligned}$$

β is given by the matrix

$$M_\beta = \begin{pmatrix} 0 & -n & 0 & -n & \dots & \dots & -n & 0 & -n & 0 & -n \\ n & 0 & 0 & 0 & \dots & \dots & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -n & \dots & \dots & -n & 0 & -n & 0 & -n \\ n & 0 & n & 0 & \dots & \dots & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \dots & \dots & -n & 0 & -n & 0 & -n \\ \vdots & \vdots & \vdots & \vdots & \ddots & & \vdots & \vdots & \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \vdots & & \ddots & \vdots & \vdots & \vdots & \vdots & \vdots \\ n & 0 & n & 0 & \dots & \dots & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \dots & \dots & 0 & 0 & -n & 0 & -n \\ n & 0 & n & 0 & \dots & \dots & 0 & n & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \dots & \dots & 0 & 0 & 0 & 0 & -n \\ n & 0 & n & 0 & \dots & \dots & 0 & n & 0 & n & 0 \end{pmatrix}$$

with respect to the basis $\varepsilon_2, \varepsilon_3, \dots, \varepsilon_{n-1}$ and its dual basis $\varphi_0(\omega_2), \varphi_0(\omega_3), \dots \varphi_0(\omega_{n-1})$.

15.5.3. $Q = E_6$. We work with the bases

$$\begin{aligned} B^+_{3,3} &= \{e_3, a_3^*a_3 - a_2^*a_2, (a_3^*a_3 - a_2^*a_2)^2, a_5^*a_5a_3^*a_3a_5^*a_5, \\ &\quad a_5^*a_5a_3^*a_3a_5^*a_5a_3^*a_3, a_3^*a_3a_5^*a_5a_3^*a_3a_5^*a_5a_3^*a_3\}, \\ B^-_{3,3} &= \{a_5^*a_5, a_3^*a_3a_5^*a_5, a_5^*a_5a_3^*a_3, a_3^*a_3a_5^*a_5a_3^*a_3, \\ &\quad a_5^*a_5a_3a_3^*(a_3^*a_3 - a_2^*a_2)^2, a_3^*a_3a_5^*a_5a_3a_3^*(a_3^*a_3 - a_2^*a_2)^2\}, \\ B^+_{6,3} &= \{a_5a_3^*a_3a_5^*a_5, a_5a_3^*a_3a_5^*a_5a_3^*a_3), \\ &\quad a_5a_3^*a_3a_5^*a_5a_3a_3^*(a_3^*a_3 - a_2^*a_2)\}, \\ B^-_{6,3} &= \{a_5, a_5a_3^*a_3, a_5a_3^*a_3(a_3^*a_3 - a_2^*a_2)\}, \\ B^+_{3,6} &= = \{a_5^*, a_3^*a_3a_5^*, (a_3^*a_3 - a_2^*a_2)a_3^*a_3a_5^*\}, \\ B^-_{3,6} &= \{a_5^*a_5a_3^*a_3a_5^*, a_3^*a_3a_5^*a_5a_3^*a_3a_5^*, a_3^*a_3a_5^*a_5(a_3^*a_3)^2a_5^*\}, \\ B^+_{6,6} &= \{e_6, a_5a_3^*a_3a_5^*a_5(a_3^*a_3)^2a_5^*\}, \\ B^-_{6,6} &= \{a_5a_3^*a_3a_5^*, a_5(a_3^*a_3)^2a_5^*\}. \end{aligned}$$

We immediately get the matrix

$$M_\beta = \begin{pmatrix} \kappa_{3,3} & \kappa_{3,6} \\ \kappa_{6,3} & \kappa_{6,6} \end{pmatrix} = \begin{pmatrix} 0 & -6 \\ 6 & 0 \end{pmatrix}$$

which represents the β with respect to the basis $\varepsilon_3, \varepsilon_6$ and dual basis $\varphi_0(\omega_3), \varphi_0(\omega_6)$.

15.5.4. E_7. For E_7 and E_8 we don't have to work with an explicit basis to calculate $\kappa_{k,l}$ since for any basis element x, $\eta(x) = \pm x$. It is enough to know the following: given any monomial $x \in e_kAe_j$ of length l, $n_{k,j}$ the number of arrows $x \in Q$ and $d(k,j)$ the distance between the vertices k, j, we know that x contains $n_{k,j} + \frac{l-d(k,j)}{2}$ arrows in Q and $d(k,j) - n_{k,j} + \frac{l-d(k,j)}{2}$ arrows in $\bar{Q}$.

We can derive the following formula:

(15.5.4)

$$\kappa_{k,j} = (-1)^{n_{k,j}} \left((d(k,j) - n_{k,j}) \frac{H_A(t)}{t^{d(k,j)}}\bigg|_{t=\sqrt{-1}} + \frac{1}{2} t \frac{d}{dt} \frac{H_A(t)}{t^{d(k,j)}}\bigg|_{t=\sqrt{-1}} \right).$$

The resulting matrix is

$$(\kappa_{k,j})_{k,j} = \begin{pmatrix} 12 & 6 & 9 & 3 & 0 & 3 & -9 \\ -6 & 0 & 3 & 0 & 0 & 0 & -3 \\ 15 & -3 & 12 & 3 & 0 & 3 & -12 \\ -3 & 0 & -3 & 0 & 0 & 0 & -6 \\ 0 & 0 & 0 & 0 & 0 & -9 & 0 \\ -3 & 0 & -3 & 0 & 9 & 0 & -6 \\ -15 & 3 & -12 & 6 & 0 & 6 & 12 \end{pmatrix}$$

A basis of $Y^*[-h-2]$ is given by

$$\varepsilon_1 = [e_1 + e_7], \varepsilon_2 = [e_2], \varepsilon_3 = [e_3 + e_7], \varepsilon_4 = [e_4], \varepsilon_5 = [e_5], \varepsilon_6 = [e_6],$$

$(\theta_0 \varepsilon_i)_{1 \le 1 \le 6}$ is given by

$$\begin{pmatrix} 3 & 6 & 0 & 3 & 0 & 3 \\ -9 & 0 & 0 & 0 & 0 & 0 \\ 3 & -3 & 0 & 3 & 0 & 3 \\ -9 & 0 & -9 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -9 \\ -9 & 0 & -9 & 0 & 9 & 0 \\ -3 & 3 & 0 & 6 & 0 & 6 \end{pmatrix} \begin{pmatrix} \varphi_0(\omega_1) \\ \varphi_0(\omega_2) \\ \varphi_0(\omega_3) \\ \varphi_0(\omega_4) \\ \varphi_0(\omega_5) \\ \varphi_0(\omega_6) \\ \varphi_0(\omega_7) \end{pmatrix}$$

Now use the defining relation of $Y[-h-2]$,

$$\varphi_0(\omega_7) = \varphi_0(\omega_1) + \varphi_0(\omega_3)$$

to obtain the matrix

$$M_\beta = \begin{pmatrix} 0 & 9 & 0 & 9 & 0 & 9 \\ -9 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 9 & 0 & 9 \\ -9 & 0 & -9 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -9 \\ -9 & 0 & -9 & 0 & 9 & 0 \end{pmatrix}$$

which represents β with respect to the basis $\varepsilon_1 \dots \varepsilon_6$ and its dual basis $\varphi_0(\omega_1), \dots, \varphi_0(\omega_6)$.

15.5.5. *E_8.* We can use 15.5.4 and get the matrix

$$M_\beta = (\kappa_{k,j})_{k,j} = \begin{pmatrix} 0 & 15 & 0 & 15 & 0 & 0 & 0 & -15 \\ -15 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 15 & 0 & 0 & 0 & -15 \\ -15 & 0 & -15 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -15 \\ 0 & 0 & 0 & 0 & 0 & 0 & -15 & 0 \\ 0 & 0 & 0 & 0 & 0 & 15 & 0 & -15 \\ 15 & 0 & 15 & 0 & 15 & 0 & 15 & 0 \end{pmatrix}$$

which represents β with respect to the basis $\varepsilon_1, \ldots \varepsilon_8$ and its dual basis $\varphi_0(\omega_1), \ldots, \varphi_0(\omega_8)$.

Remark 15.5.5. With respect to our chosen bases $(\varepsilon_i)_{i \in I'}$ and $\phi_0(\omega_i)_{i \in I'}$, such that the vertex set $I' \subset I$, together with the arrows in I form a connected subquiver $\bar{Q}'$, M_β can be written in this general form:

$$M_\beta = \frac{h}{2} \cdot (C')^\epsilon, \tag{15.5.6}$$

where we call $(C')^\epsilon$ the *signed adjacency matrix* of the subquiver $\bar{Q}'$, that is

$$(C')_{ij} = \begin{cases} 0 & \text{if } i, j \text{ are not adjacent,} \\ +1 & \text{if arrow } i \leftarrow j \text{ lies in } Q^*, \\ -1 & \text{if arrow } i \leftarrow j \text{ lies in } Q, \end{cases} \tag{15.5.7}$$

In the D_{n+1}-case, we have

$$M_\beta = n \cdot \begin{pmatrix} 0 & 1 & 0 & \cdots & \cdots & \cdots & \cdots & 0 \\ -1 & 0 & 1 & 0 & \cdots & \cdots & \cdots & 0 \\ 0 & -1 & 0 & 1 & 0 & \cdots & \cdots & 0 \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots & & \vdots \\ \vdots & & \ddots & \ddots & \ddots & \ddots & \ddots & \vdots \\ \vdots & & & \ddots & \ddots & \ddots & \ddots & 0 \\ \vdots & & & & \ddots & \ddots & \ddots & 1 \\ 0 & \cdots & \cdots & \cdots & \cdots & 0 & -1 & 0 \end{pmatrix}^{-1},$$

in the E_6-case, we have

$$M_\beta = 6 \cdot \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}^{-1},$$

in the E_7-case, we have

$$M_\beta = 9 \cdot \begin{pmatrix} 0 & -1 & 0 & 0 & 0 & 0 \\ 1 & 0 & -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 & 0 & 1 \\ 0 & 0 & 0 & 0 & -1 & 0 \end{pmatrix}^{-1},$$

and in the E_8-case, we have

$$M_\beta = 15 \cdot \begin{pmatrix} 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 & -1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 \end{pmatrix}^{-1}.$$

16. PRODUCTS INVOLVING $HH^2(A)$

We start with $HH^2(A) \times HH^3(A) \to HH^5(A)$ first and then deduce $HH^2(A) \times HH^2(A) \to HH^4(A)$ from associativity.

16.1. $HH^2(A) \times HH^3(A) \to HH^5(A)$.

We will prove the following general proposition:

Proposition 16.1.1. *For the basis elements* $f_i \in HH^2(A)$, $h_j \in HH^3(A)$, *the cup product is*

$$f_i h_j = \delta_{ij}\psi_0. \tag{16.1.2}$$

Proof. Recall the maps

$$\begin{array}{rcl} h_j : A \otimes \mathcal{N} & \to & A, \\ 1 \otimes 1 & \mapsto & \omega_j \end{array}$$

and lift it to

$$\begin{array}{rcl} \hat{h}_j : A \otimes \mathcal{N} & \to & A \otimes A, \\ 1 \otimes 1 & \mapsto & 1 \otimes \omega_j. \end{array}$$

Then

$$\hat{h}_j(d_4(1 \otimes a \otimes 1)) = \hat{h}_j(a \otimes 1 - 1 \otimes a) = a \otimes \omega_j = d_1(1 \otimes a \otimes \omega_j),$$

so

$$\begin{array}{rcl} \Omega h_j : \Omega^4(A) & \to & \Omega(A), \\ 1 \otimes a \otimes 1 & \mapsto & 1 \otimes a \otimes \omega_j. \end{array}$$

Then we have

$$\begin{array}{rcl} \Omega h_j(d_5(1 \otimes 1)) & = & \Omega h_j(\displaystyle\sum_{a \in \bar{Q}Q} \epsilon_a a \otimes a^* \otimes 1 + \sum_{a \in \bar{Q}} \epsilon_a 1 \otimes a \otimes a^*) \\ & = & \displaystyle\sum_{a \in \bar{Q}} \epsilon_a \otimes a^* \otimes \omega_j = d_2(1 \otimes \omega_j), \end{array}$$

so

$$\begin{array}{rcl} \Omega^2 h_j : \Omega^5(A) & \to & \Omega^2(A), \\ 1 \otimes 1 & \mapsto & 1 \otimes \omega_j. \end{array}$$

This gives us

$$f_i(\Omega^2 h_j)(1 \otimes 1) = f_i(1 \otimes \omega_j) = \delta_{ij}\omega_j,$$

i.e. the cup product

$$f_i h_j = \delta_{ij}[\omega_j] = \delta_{ij}\psi_0.$$

□

16.2. $HH^2(A)\times HH^2(A)\to HH^4(A)$. Since $\deg HH^2(A)=-2$, their product has degree -4 (i.e. lies in $span(\zeta_0)$, so it can be written as

$$\begin{array}{rcl} HH^2(A)\times HH^2(A) & \to & HH^4(A), \\ (a,b) & \mapsto & \langle -,-\rangle\zeta_0, \end{array}$$

where $\langle -,-\rangle : HH^2(A)\times HH^2(A)\to\mathbb{C}$ is a bilinear form. We prove the following proposition:

Proposition 16.2.1. *The cup product $HH^2(A)\times HH^2(A)\to HH^4(A)$ is given by $\langle -,-\rangle=\alpha$, where α (from Proposition 15.2.3) is regarded as a syymetric bilinear form.*

Proof. We use (15.4.2) to get

$$\theta_0(f_if_j)=\theta_0(\langle f_i,f_j\rangle\zeta_0)=\langle f_i,f_j\rangle\psi_0. \tag{16.2.2}$$

On the other hand, by Proposition 15.2.3 and Proposition 16.1.1,

$$(\theta_0f_i)f_j=\alpha(f_i)f_j=\sum(M_\alpha)_{li}h_lf_j=(M_\alpha)_{ji}\psi_0=(M_\alpha)_{ij}\psi_0. \tag{16.2.3}$$

By associativity of the cup product, we can equate (16.2.2) and (16.2.3) to get

$$\langle f_i,f_j\rangle=(M_\alpha)_{ij}. \tag{16.2.4}$$

□

16.3. $HH^2(A) \times HH^4(A) \xrightarrow{0} HH^6(A)$**.** This computation uses the Batalin-Vilkovisky structure on Hochschild cohomology: We have $\deg HH^2(A) = -2$, $\deg HH^4(A) \geq -h$ and $\deg HH^6(A) \leq -h-2$. So we know by degree argument that

$$f_k\zeta_l = \begin{cases} 0 & l > h-4 \\ \sum\limits_s \lambda_s \varphi(\omega_s) & l = h-4 \end{cases} . \tag{16.3.1}$$

We use [Eu3, (6.0.12)] and the isomorphism $HH^i(A) = HH_{6m+2-i}(A)$ to get for the Gerstenhaber bracket on $HH^*(A)$:

$$\begin{aligned} [f_k, \zeta_l] &= \Delta(f_k\zeta_l) - \underbrace{\Delta(f_k)}= 0\zeta_l - f_k \underbrace{\Delta(\zeta_l)}_{=0} \\ &= \sum_s \lambda_s(\frac{1}{2}+m)h\beta^{-1}(\varphi(\omega_s)) \end{aligned}$$

The Gerstenhaber bracket has to be independent of the choice of $m \geq 0$. This implies that the RHS has to be zero, so all $\lambda_s = 0$. This shows that

$$f_k\zeta_{h-4} = 0, \tag{16.3.2}$$

so we have that the cup product of $HH^2(A)$ with $HH^4(A)$ is zero.

16.4. $HH^2(A) \times HH^5(A) \xrightarrow{0} HH^7(A)$**.** Let $a \in HH^2(A)$ and $b \in HH^5(A)$ be homogeneous elements, then $ab = \lambda\theta_k \in HH^7(A) = U[-2h-2]$, $\lambda \in \mathbb{C}$. Then

$$\lambda\psi_0 = \lambda\psi_k z_k = \lambda\theta_k\zeta_k = \lambda b(a\zeta_k) = 0,$$

the last equality coming from the product $a\zeta_k \in HH^2(A) \cup HH^4(A) = 0$.

17. Products involving $HH^3(A)$

17.1. $HH^3(A) \times HH^3(A) \xrightarrow{0} HH^6(A)$**.** This follows by degree argument: $\deg HH^3(A) = -2$, $\deg HH^6(A) \leq -h-2 < -4$.

17.2. $HH^3(A) \times HH^4(A) \xrightarrow{0} HH^7(A)$**.** This follows by degree argument: $\deg HH^3(A) = -2$, $\deg HH^4(A) \geq -h$, $\deg HH^7(A) \leq -h-4 < -h-2$.

17.3. $HH^3(A) \times HH^5(A) \xrightarrow{0} HH^8(A)$**.** This follows by degree argument: $\deg HH^3(A) = -2$, $\deg HH^5(A) \geq -h-2$, $\deg HH^8(A) = -2h-2 < -h-4$.

18. Products involving $HH^4(A)$

18.1. $HH^4(A) \times HH^4(A) \xrightarrow{0} HH^8(A)$**.** This follows by degree argument: $\deg HH^4(A) \geq -h$, $\deg HH^8(A) = -2h-2 < -2h$.

18.2. $HH^4(A) \times HH^5(A) \xrightarrow{0} HH^9(A)$**.** This is clear for $Q = D_{n+1}$, n odd, $Q = E_7$, E_8 where $HH^9(A) = K[-2h-2] = 0$.

Let $Q = D_{n+1}$, n even or $Q = E_6$. Let $a \in HH^4(A)$, $b \in HH^5(A)$.The product $HH^2(A) \times HH^3(A) \to HH^5(A)$, $(x, y) \mapsto \langle x, y\rangle\zeta_0$ induces a nondegenerate bilinear form $\langle -, -\rangle$. If $ab \in HH^9(A) = HH^3(A)[-2h]$ is nonzero, then we can find a $c \in HH^2(A)$, such that $c(ab) = \zeta_0$. But this equals $(ca)b = 0$ since $HH^2(A) \times HH^4(A) \xrightarrow{0}$ which gives us a contradiction.

19. $HH^5(A) \times HH^5(A) \to HH^{10}(A)$

Proposition 19.0.1. *The multiplication of the subspace* $U[-2]^*$ *with* $HH^5(A)$ *is zero.*

The pairing on $Y^*[-h-2]$ *is*

$$\begin{array}{rcl} Y^*[-h-2] \times Y^*[-h-2] & \to & HH^{10}(A), \\ (a, b) & \mapsto & \Omega(a, b)\varphi_4(\zeta_0), \end{array} \tag{19.0.2}$$

where the skew-symmetric bilinear form $\Omega(-, -)$ *is given by the matrix* $-M_\beta$ *from subsection 15.5.*

Proof. We have $\deg HH^5(A) \geq -h-2$ and $\deg HH^{10}(A) \leq -2h-4$, so we can get a nonzero multiplication only by pairing bottom degree parts of $HH^5(A)$ which is $Y^*[-h-2]$. The product lies in the top degree part of $HH^{10}(A) = HH^4(A)[-2h]$ which is spanned by $\varphi_4(\zeta_0)$. This gives us the pairing of the form (19.0.2).

We want to find the matrix $(\Omega(\varepsilon_i, \varepsilon_j))_{i,j}$ where ε_i are a basis of $Y^*[-h-2]$, given in the section about $HH^5(A)$. Recall that the multiplication $HH^1(A) \times HH^5(A) \to HH^6(A)$ was given by a skew-symmetric matrix $((M_\beta)_{i,j})_{i,j\in F}$, so that $\theta_0\varepsilon_i = \sum\limits_{k\in F} (M_\beta)_{k,i}\varphi_0(\omega_k)$.

We multiply $\varepsilon_i\varepsilon_j = \Omega(\varepsilon_i, \varepsilon_j)\varphi_4(\zeta_0)$ with θ_0 (see 15.4.2):

$$\theta_0(\varepsilon_i\varepsilon_j) = \Omega(\varepsilon_i, \varepsilon_j)\varphi_5(\psi_0). \tag{19.0.3}$$

Using associativity, this equals

$$(\theta_0\varepsilon_i)\varepsilon_j = \sum_{k\in F} (M_\beta)_{k,i}\varphi_0(\omega_k)\varepsilon_j = (M_\beta)_{j,i}\psi_0 = -(M_\beta)_{i,j}\varphi_5(\psi_0). \tag{19.0.4}$$

We see from equations (19.0.3) and (19.0.4) that

$$\Omega(\varepsilon_i, \varepsilon_j) = -(M_\beta)_{i,j}.$$

□

References

[CBEG] W. Crawley-Boevey, P. Etingof, V. Ginzburg, *Noncommutative Geometry and Quiver algebras,* `arXiv:math.AG/0502301`.

[CST] J. Cuntz, G. Skandalis and B. Tsygan: *Cyclic homology in non-commutative geometry*, Encyclopedia of Mathematical Sciences, 121. Operator Algebras abd Non-Commutative Geometriy, II. Springer-Verlang, Berlin, 2004.

[D] V.A. Dolgushev, A Proof of Tsygan's formality conjecture for an arbitrary smooth manifold, PhD thesis, MIT; `arxiv:math.QA/0504420`.

[DGT] Yu. Daletski, I. Gelfand and B. Tsygan, *On a variant of noncommutative geometry*, Soviet Math. Dokl. **40**, 2 (1990) 422-426.

[EE2] P. Etingof and C.Eu: *Hochschild and cyclic homology of preprojective algebras of ADE quivers* `arXiv:math.AG/0609006`

[ES1] K. Erdmann, N. Snashall: Preprojective algebras of Dynkin type, periodicity and the second Hochschild cohomology. Algebras and modules, II (Geiranger, 1996), 183–193, CMS Conf. Proc., 24, Amer. Math. Soc., Providence, RI, 1998.

[ES2] K. Erdmann, N. Snashall, On Hochschild cohomology of preprojective algebras. I, II. J. Algebra 205 (1998), no. 2, 391–412, 413–434.

[Eu3] C. Eu, *The calculus structure of the Hochschild homology/cohomology of preprojective algebras of Dynkin quivers*, `arXiv:0706.2418v1`

[MOV] A. Malkin, V. Ostrik and M. Vybornov: *Quiver varieties and Lusztig's algebra* `arXiv:math.AG/0403222`

[S] A. Schofield: *Wild algebras with periodic Auslander-Reiten translate*, preprint.

Department of Mathematics, Massachusetts Institute of Technology, Cambridge, MA 02139, U.S.A.

E-mail address: `ceu@math.mit.edu`